



SUPPLEMENTAL LABELING

EVITO® T Fungicide For Use on Wheat and Sweet Corn

GROUP	11	FUNGICIDE
GROUP	03	FUNGICIDE

ACTIVE INGREDIENT:

Fluoxastrobin: [(1E)-[2-[[6-(2-Chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl] (5,6-dihydro-1,4,2-dioxazin-3-yl) methanone-O-methyloxime]	18%
Tebuconazole: alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol	25%
OTHER INGREDIENTS:	57%
TOTAL:	100%

This product contains **1.67** lbs of fluoxastrobin and **2.32** lbs tebuconazole per gallon

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label find someone to explain it to you in detail.)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, worker protection standard requirements, and precautions on the EPA registered main label EVITO® T Fungicide (EPA Reg. No. 66330-383).

This Supplemental Label must be in the possession of the user at the time of application.

This Supplemental Label expires on May 16, 2014.

EPA Reg. No. 66330-383

Wheat

Disease Control	Rate to Use	Application Timing and Resistance Management
<p>Leaf rust (<i>Puccinia recondita f. sp. tritici</i>)</p> <p>Stripe rust (<i>Puccinia striiformis</i>)</p> <p>Stem rust (<i>Puccinia graminis</i>)</p> <p>Septoria leaf and glume blotch (<i>Septoria tritici</i>, <i>Septoria nodorum</i>)</p> <p>Tan Spot (<i>Pyrenophora tritici-repentis</i>)</p>	<p>4 - 6 fl oz/A*</p>	<ul style="list-style-type: none"> • Apply a maximum of two applications per season. • Apply no later than Feekes growth stage 10.5. • For optimum results, apply the first application at approximately Feekes growth stage 5 (Zadoks 31)(shooting-pseudostem erected) and a second application no later than Feekes growth stage 10.5 (Zadoks 54)(heading completed).

*(0.052 lbs fluoxastrobin and 0.072 lbs tebuconazole per acre to 0.078 lbs fluoxastrobin and 0.181 lbs tebuconazole per acre.)

RESTRICTIONS AND OTHER INFORMATION:

- Do not apply more than 12 fl oz (0.156 lbs fluoxastrobin+ 0.037 lbs tebuconazole.) of EVITO T Fungicide per acre per crop season.
- There is a maximum of two applications per crop season.
- Do not apply EVITO T Fungicide after Feekes growth stage 10.5 (heading completed).
- Restricted-entry interval (REI) = 12 hours.
- Apply in a minimum of 10 gallons of water per acre by ground and 3 gallons of water per acre by air.
- Do not apply EVITO T Fungicide within 40 days of harvest for grain or straw.
- Do not apply EVITO T Fungicide within 7 days of harvest for forage or hay.
- Do not allow livestock to graze or feed on green forage within 7 days after treatment with EVITO T Fungicide.

SWEET CORN

Disease Control	Rate to Use	Application Timing and Resistance Management
Rust, common <i>(Puccinia sorghi)</i> Rust, southern <i>(Puccinia polyspora)</i> Anthracnose leaf blight <i>(Colletotrichum graminicola)</i> Gray Leaf Spot <i>(Cercospora sorghi)</i> Northern corn leaf blight <i>(Setosphaeria turcica)</i> Northern corn leaf spot <i>(Cochliobolus carbonum)</i> Southern corn leaf blight <i>(Cochliobolus heterostrophus)</i> Eye Spot <i>(Aureobasidium zeae)</i>	4-9 fl oz/A*	<ul style="list-style-type: none"> • Begin applications preventatively. Apply as needed on a 14 day interval. <p>Resistance Management:</p> <p>To limit the potential for development of disease resistance:</p> <ul style="list-style-type: none"> • In areas with typically 1-4 sprays per year, alternate every application of EVITO T Fungicide with at least one application of another effective mode of action fungicide. • Do not make more than two (2) sequential applications of EVITO T Fungicide before alternating with a different mode of action for at least one (1) application

*0.052 lbs fluoxastrobin and 0.072 lbs tebuconazole per acre to 0.117 lbs fluoxastrobin and 0.163 lbs tebuconazole per acre.

RESTRICTIONS AND OTHER INFORMATION:

- Do not apply more than 36 fl oz (0.468 lbs fluoxastrobin+0.652 lbs tebuconazole) of EVITO T Fungicide per acre per year.
- There is a maximum of two applications per crop season, and a minimum retreatment interval of 14 days.
- Do not apply EVITO T Fungicide within 7 days of harvest of ears or for forage.
- Restricted-entry interval (REI) = 19 days.
- Do not apply EVITO T Fungicide within 49 days of harvest for fodder.
- Apply in a minimum of 10 gallons of water per acre by ground and 3 gallons of water per acre by air.

Produced for:

ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513



EVITO is a registered trademark of Arysta LifeScience North America, LLC
 Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience

Supplemental label approved EPA May 12, 2011