



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9615

**FOR DISTRIBUTION AND USE
IN ALL STATES EXCEPT TEXAS**

**THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS MADE
AS PERMITTED UNDER FIFRA SECTION 2(ee)
AND HAS NOT BEEN SUBMITTED TO OR
APPROVED BY THE US EPA.**

This label for EXIREL® insect control is valid until December 31, 2024, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR CONTROL OF CARROT WEEVIL IN FLORENCE FENNEL IN ALL STATES EXCEPT TEXAS

Directions For Use

EXIREL insect control is recommended for control of carrot weevil in Florence Fennel.

Apply EXIREL insect control at 13.5-20.5 fl oz/A as a foliar spray using properly calibrated ground spray equipment. Initial application should be made at first sign of adult weevil presence.

For best results use the highest labeled rate of EXIREL insect control (20.5 fl oz/A). Under moderate to high pest pressure, an additional application of EXIREL insect control may be needed to maintain weevil populations below damaging levels. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for this pest.

Thorough spray coverage of is essential to achieve best results.

Read and follow the protection of pollinators section of the federal label.

FMC will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by FMC. User assumes all risks associated with uses not specified by the label or FMC 2(ee) instructions.

RESTRICTIONS

Do not apply a total of more than 0.4 lbs ai/A of CYAZYPYR® active or cyantraniliprole containing products per calendar year.

The minimum retreatment interval is 5 days.

The pre-harvest interval (PHI) is 1 day.

The field re-entry interval is 12 hours.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104

IMPORTANT NOTICE

BEFORE BUYING OR USING FMC PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced FMC product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the FMC product(s) covered by this Bulletin. FMC will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by FMC on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.**

R-4353 011520 01-14-20

FMC, EXIREL, and CYAZYPYR are trademarks of FMC Corporation or an affiliate.
©2020 FMC Corporation



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104