



YaraVita™

**Ferrichel
Ninety**™



**Net weight
10lb/
4.5 kg**

NOTICE

Before using this product, read the entire label

Ferrichel Ninety™

12-13-0

A formulated soluble powder iron with nitrogen and phosphorus

Guaranteed analysis:

Total Nitrogen (N)	12%
12% Ammoniacal Nitrogen	
Available Phosphate (P₂O₅)	13%
Iron (Fe)	9%

 9% chelated iron

Derived from Ammonium Phosphate, Ammonium Sulfate, Ferric Citrate

F1673

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

This product is recommended for use in conjunction with a complete fertilization program based on soil or tissue analysis.

Contents made in Great Britain

Guaranteed & Manufactured by Yara UK Ltd., Manor Place, Wellington Road, The Industrial Estate, Pocklington, York YO42 1DN, U.K.
Manufactured for Yara North America, Inc. 100 North Tampa Street, Suite 3200, Tampa, FL 33602 Customer Service: 800-234-9376

Warning

Contains ammonium iron(III) citrate, ammonium sulphate.

H319:Causes serious eye irritation.

H315:Causes skin irritation.

P280:Wear protective gloves and eye protection.

P264:Wash hands thoroughly after handling.

P305:IF IN EYES:

P351:Rinse cautiously with water for several minutes.

P338:Remove contact lenses, if present and

easy to do. Continue rinsing.

P337:If eye irritation persists:

P313-a:Get medical attention.

P302:IF ON SKIN:

P352:Wash with plenty of soap and water.

P332:If skin irritation occurs:

P313:Get medical attention.



11.04.19

WLU5/451

YaraVita™ Ferrichel Ninety™

APPLICATION RATES AND TIMINGS

Apples, Pears: 1 to 3 lb/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering.

Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.

Apricots, Cherries, Nectarines, Peaches, Plums: 1 to 3 lb/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre

Citrus: 1 to 3 lb/acre during Spring and Autumn flushes, at 10 to 14 day intervals as necessary. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.

Conifers: 1 to 3 pounds/acre at start of new season leaf production and again in early autumn. Water rate: 50 to 100 gallons/acre.

Cucurbits (Field Grown): 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.

Lettuce (field grown): 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.

Nuts (deciduous): 1 to 3 lbs/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.

Melon, Water Melons (Field Grown): 1 to 3 lbs/acre from two weeks after transplanting, or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre

Onions: 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.

Peppers, Tomatoes (Field Grown): Peppers (field grown): 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.

Soybeans: 1 to 3 lbs/acre from 2 weeks after emergence, or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Do not apply during flowering. Water rate: 20 gallons/acre.

Sugar Cane: 1-3 lb/acre when plants are 12" tall and again 10 to 14 days later. Water rate: 20 gallons/acre.

Sweet Corn: 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.

Vines: 1 to 3 lb/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.

Extended recommendations available at

<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and Reproductive Harm –www.P65Warnings.ca.gov