

SACCO		FERRILENE 20 LBS USA		EN		4105867F07P1S9 04	
STF_PPCOEXPETMETPE - TRIPLEX_70				MULTI FONT			
830x540	3	K	P.2062				
2	K	P.2062					



002574

FERRILENE

20Lb 9.07 kg NET WEIGHT

SOLUBLE MICRONUTRIENT GRANULE

GUARANTEED ANALYSIS

Iron (Fe) 6.0 %
 6.0 % Chelated Iron (Fe)
 Derived from: Iron EDDHA

GENERAL INFORMATION

FERRILENE is a source of Iron Chelate readily available for crops, especially in alkaline soil conditions. The product is to be applied into the root zone through drip line, drench, injection, band application (watering in the solution). Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn/garden.

NOTICE

Direct exposure to sunlight is to be avoided.

KEEP OUT OF REACH OF CHILDREN

Information regarding the contents and levels of metals in this product is available on the Internet at www.aapfc.org/metals.htm. Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rains are expected.

STORAGE

Keep the product sheltered from the sun and in a dry warehouse. Once open, completely use the contents of the bag. When applied do not leave exposed to sunlight and cover immediately with soil.


CLEANING OF SPRAYER

Flush with clean water all parts of the sprayer including the pump, hoses and nozzles several times prior to and after the use of this product.

FERRILENE is a Registered Trademark of VALAGRO S.p.A.

Distributed and guaranteed by Valagro USA Inc., 2020 Ponce de Leon Blvd., Ste. 1203
 Coral Gables, FL 33134 - Phone +1-786-230-1020 / +1-786-230-1019 / Fax +1-786-475-9032

ROOT APPLICATION



FERRILENE | Valagro®

FERRILENE | Valagro®



www.valagro.com

FERRILENE

FIELD APPLICATION INSTRUCTIONS

The product can be applied throughout the season, but applications during the early stages of vegetative development or post harvest (before dormancy) are preferred. Splitting smaller amounts of product and making several applications, is preferable to applying large quantities of Ferrilene in fewer applications. Rates depend on deficiency, severity and soil conditions (lower rates with more applications on sandy soils).

CROPS	RATE	TIMING
Fruit and Nuts crops: apples, apricots, avocados, cherries, mandarin, peaches, pistachio, almonds, nectarines, peaches, plums, plums, pines and walnuts	5-10 lb/Ac per application.	Apply annually in the spring, either prior to or at the time of new spring growth. Post harvest application prior dormancy are other recommended.
Citrus	5-15 lb/Ac per application.	Broadcast evenly within the drip line prior to the winter flush of growth.
Grapes	5-10 lb/Ac per application.	Early spring or when deficiency symptoms first appear.
Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries, Blackberries, and Strawberries	10-20 lb/Ac per application.	Early spring or when deficiency symptoms first appear.
Field crops: Alfalfa and forage grasses, Cotton, Corn, field peas, flax, rice, soybeans, sorghum, Sugar beets, and sugar cane	2-10 lb/Ac per application.	At planting or when deficiency symptoms first appear.
Corn, Peanuts, and sorghum	Band or side dress: 2-10 lb/Ac.	At planting or when deficiency symptoms first appear.
Vegetable crops: beans, corn, mustard and spinach	Band or side dress: 5-10 lb/Ac.	At planting or when deficiency symptoms first appear.
Black-eyed peas, Carrots, Cucumbers, eggplants, Melons, Onions, Peas, Peas, Peppers, Potatoes, Radishes, Squash, Tomatoes, and Turnips	Band or side dress: 5-10 lb/Ac.	At planting or when deficiency symptoms first appear.
Cabbage, Cauliflower, Celery, and lettuce	Band or side dress: 5-10 lb/Ac.	At planting or when deficiency symptoms first appear.
Ornamentals	Band or side dress: 5-10 lb/Ac.	At planting or when deficiency symptoms first appear.
Turf	2-3 lb/1000 sq ft when deficiency symptoms appear. 10-15 lb/1000 sq ft maintenance.	At planting or when deficiency symptoms first appear.

Use lower rates in maintenance situations and higher rates in deficiency situations. The actual rate applied should be based on soil type and severity of iron deficiency.

CONDITIONS OF SALE

Seller warrants this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with direction under normal condition and use. No one other than an office of the seller is authorized to make any warranty, guaranty or direction concerning this product. Because time, place, rate of application and other conditions of use are beyond the seller control, the seller's liability from handling, storage, and use of this product is limited to replacement of the product or a refund of the purchase price. It is impossible to eliminate all risks associated with the use of this product. Crop injury, lack of performance, all other unintended consequences may result because of such factors as use of the product contrary to label instructions, abnormal condition, presence of other material, the manner of application, or other factors, all of which are beyond the control of Valagro USA or the seller. All such risks shall be assumed by buyer.

GREENHOUSE APPLICATION INSTRUCTIONS: DO NOT

exceed 10 ounces per 25 gallons for greenhouse crops. Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rains are expected.



4105867F07P1S9 04