GENERAL INFORMATION

Foli-Zyme is formulated as a soluble concentrate that requires a minimum 1 to 10 dilution with water for all applications. **Foli-Zyme** is a fluid nutrient compound designed to prevent nutrient deficiencies in plants.

DIRECTIONS FOR USE

Recommended use:

Foli-Zyme may be used on the following crops at recommended rates and application intervals below: alfalfa, almonds, apples, artichokes, asparagus, avocados, barley, all beans, broccoli, cabbage, carrots, cauliflower, celery, citrus, corn, cotton, grapes, hops, lettuce, melons, milo, nectarines, peaches, peanuts, peas, pecans, peppers, pistachios, potatoes, rice, soybean, sorghum, strawberries, sugar beets, sugar cane, sunflowers, sweet corn, tobacco, tomatoes, turnips, walnuts, watermelons, wheat and most other crops.

Field row crops and vegetables: Foliar spray **Foli-Zyme** on the following schedule: Apply 1 gallon per acre (9 liters per hectare). Start applications when the crop has 2 to 3 leaves or is 20 days old. Continue sprays at 10 to 14 day intervals up to 6 times annually.

Tree, fruits and nuts: Apply 1 gallon per acre (9 liters per hectare) up to 6 times annually, beginning with delayed dormant spray. Utilize foliar application interval of 14 days. AVOID PERIODS WHEN FRUIT IS EXTREMELY SENSITIVE TO SPRAY DAMAGE, PRIMARILY DURING THE POLLINATION PERIOD.

Citrus: Apply 1 gallon per acre (9 liters per hectare) every 30 days. Do not mix with petroleum spray oils or ammoniated copper compounds.

Solid-seeded field crops: Apply 1 gallon per acre (9 liters per hectare) when the crop is 3 inches high. Continue foliar application on 14-day interval until crop in 14 days from harvest.

Mixing Instructions:

Many pesticides can tank mixed with **Foli-Zyme**. Follow this mixing procedure: 1. Water, 2. **Foli-Zyme**, 3. Pesticide **Foli-Zyme** will disperse in water with little agitation.

CAUTION: Be sure to conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility. Foli-Zyme has a pH range of 2.0 to 3.5 and will buffer spray tank water. The use of additional buffering agents may result in crop damage.

WARNING: Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat, and/or skin. Avoid contact with eyes, skin and clothing.

IF SWALLOWED: Induce vomiting be sticking finger down throat. Never give anything by mouth to an unconscious person. Get medical attention

IF IN EYES: Wash eyes with plenty of clear water for at least 15 minutes. Get medical attention.

Notice: The information contained herein is true and accurate to the best of our knowledge and belief, but is presented without guarantee since field conditions and use are beyond our control. Neither the manufacturer nor the seller make any warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instruction. Read and follow the label directions carefully.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee of direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond sellers control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.



GUARANTEED ANALYSIS

Total Nitrogen (N)	15.0%
Soluble Potash (K,O)	
Calcium (Ca)	

Derived from urea, potassium chloride, and calcium chloride.

CAUTION

KEEP OUT OF REACH OF CHILDREN; MAY BE HARMFUL IF SWALLOWED, INHALED OR IN CONTACT WITH SKIN; MAY CAUSE EYE IRRITATION. MAY CAUSE DAMAGE THROUGH PROLONGED EXPOSURE.

PRECAUTIONS: Do not breathe dust/fume/gas/mist/vapors/spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in well-ventilated area. Avoid release to environment. Wear protective gloves/protective clothing/eye protection/face protection. Keep container tightly closed. Keep cool and protected from sunlight.

If medical advice is needed, have product container or label at hand. For more details regarding the safe use/handling, storage and disposal of this product as well as additional hazards and first aid measures, please obtain the current Safety Data Sheet at www.stollerusa.com.

STORAGE AND DISPOSAL

We recommend storing pails a maximum of three pails high. Do not stack pallets. Store container tightly closed in a cool/well-ventilated dry place protected from sunlight above 40°F. Do not allow container to freeze. Store locked up. DISPOSE of contents/ container at a permitted facility or as advised by your local regulatory authority. Do not contaminate water, food or feed by storage or disposal. Do not apply directly to areas where surface water is present, or to intertidal areas below the mean high water mark.Do not contaminate water when disposing of equipment wash water. Do not reuse empty container.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html

Foli-Zyme

Foliar Nutrient Compound with 3.0% Calcium (Contains N + Calcium)

15-0-2

2399

Density: 10.70 lb/gal (1.27 kg/L) @ 68°F (20°C)

NET CONTENT NET WEIGHT

 □ 2.50 gal (9.5 L)
 26.8 lb (12.0 kg)

 □ 2.64 gal (10.0 L)
 28.2 lb (12.7 kg)

 □ 5.00 gal (18.9 L)
 53.5 lb (24.0 kg)

 □ 5.28 gal (20.0 L)
 56.5 lb (25.4 kg)

 □ 55.00 gal (208.2 L)
 588.5 lb (264.4 kg)

 □ 275.00 gal (1041.0 L)
 2942.5 lb (1322.1 kg)

Z- FZ 15-0-2 Rev: 20180817

Manufactured and Guaranteed by Stoller Enterprises, Inc.

9090 Katy Freeway, Suite 400 Houston, TX 77024 Phone: 1-800-539-5283 or 713-461-1493



