



FOLIAR PHITE

FERTILIZERS



DIRECTIONS FOR USE

Vegetable Crops: Apply 2 to 4 pints per acre with sufficient water for thorough coverage. Apply before flowering on such crops as corn or beans. On cole crops, apply at 3 to 5 true leaf stage. On peas and lentils, apply at second true leaf. Apply as required several weeks apart.

Cucurbits, Melons, Yams, Potatoes: Apply 1 to 3 pints per acre with sufficient water for thorough coverage. Apply prior to each major bloom. Avoid high temperature, low humidity conditions in combination with such products as dimethoate. If in doubt, treat a small area of the field before large-scale application.

Cereal Crops: Apply 1 to 3 pints per acre with sufficient water for thorough coverage. Apply prior to bloom.

Field and Row Crops: Apply 1 to 3 pints per acre with sufficient water for thorough coverage. Make application prior to bloom on a crop such as cotton, or within 40 days of a plant's development on a crop such as corn.

Deciduous Tree Crops: Apply 2 to 4 pints per acre with sufficient water for thorough coverage. Apply before bloom. Repeat at petal fall. Continue applications as required.

Citrus: Apply 1 to 4 pints per acre with sufficient water for thorough coverage. Apply before fall and spring new growth. Apply before bloom.

Vines (Table, Wine, Raisin, or other Berries): Apply 2 to 4 pints per acre several weeks before bloom with sufficient water for thorough coverage. Do not overload spray mixture with other high salt fertilizers and pesticide mixtures. Apply 1 to 2 pints per acre with sufficient water for thorough coverage after shatter. Apply 1 to 2 pints per acre with sufficient water for thorough coverage several weeks before harvest. **Soil:** Apply 1 to 4 pints per acre with irrigation water or liquid fertilizer. Please check compatibility before large-scale use.

Hops: Apply up to 3 quarts per acre at 6 to 12-inch shoot growth. Make additional applications after training (when vines are 5 to 6 feet tall), approximately 3 weeks later. Repeat application 1 to 2 times with an interval of 10 to 15 days or as needed.

Onion, Garlic and other Vegetable Root, Bulb or Tuber Crops: Apply 1 to 3 pints per acre when plants are well established. Repeat application 1 to 2 times with an interval of 10 to 15 days or as needed.

0-0-28

GUARANTEED ANALYSIS

Soluble Potash (K_2O) 28.00%

DERIVED FROM: Potassium Hydroxide.

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS:

Phosphorous Acid..... 35.78%

KEEP OUT OF REACH OF CHILDREN



WARNING

Harmful if swallowed. May cause respiratory irritation.

PRECAUTIONARY STATEMENTS

Avoid breathing mist or vapor. Wash thoroughly after handling.

Do not eat, drink or smoke when using this product.

Use only outdoors or in a well-ventilated area.

FIRST AID

In all cases, call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Call a poison control center or doctor if you feel unwell.

Rinse mouth. **IF INHALED:** Remove person to fresh air and keep comfortable for breathing. Call a poison control center or doctor if you feel unwell.

Weight Per Gallon: 12.8 lb., pH 6.2

Crystallization Point: -10°F (Will resolublize if shaken).

NET CONTENTS _____ GALLONS

GENERAL INFORMATION

FoliarPhite is a highly concentrated foliar and soil phosphite fertilizer. It is compatible with most pesticides (See PRECAUTIONS). A jar test to ensure compatibility before performing any large-scale mixing is recommended.

USAGE PRECAUTIONS

FoliarPhite is generally compatible with most pesticides. This product has a pH of 6.2 which may result in less buffer or acidifying agent being required to adjust the pH to the desired level. Always add **FoliarPhite** to the spray tank first. Use surfactants at rates sufficient to wet the plants. Do not use excessive rates.

Do not use in combination with excessive pesticide and nutrient tank mixes, lime lignosulfonates (not compatible), or high rates of metals (sulfates, oxides, or chlorides). Application of this product with several emulsifiable concentrates during hot, dry conditions with insufficient water may cause margin leaf burn. Under these conditions use sufficient water for thorough coverage. Applications made with this product in high concentrates and in combination with known phytotoxic materials should be avoided or test the combination on a small scale prior to application on a large scale. If margin burn or scarring is evident, do not use or wait for cooler temperatures. Do not apply with products known to be phytotoxic with phosphite fertilizers such as copper products (cupric oxides, cupric hydroxides, lime-copper sulfate mixtures). When in doubt of a tank mix phytotoxicity, test a small area before large-scale spraying. For commercial use only. Do not use in green houses or on ornamental crops such as flowers or shrubs.

NOTICE: Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfco.org/metals.html>.

STORAGE AND DISPOSAL

Store in a well-ventilated place and keep locked up. Store in original container only and keep tightly sealed when not in use. Store product and dispose of unused product and empty containers in accordance with federal, state and local regulations.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the face hereof. All statements made concerning this product apply only when used as directed.

Manufactured for:
GENESIS AGRI • PRODUCTS INC.
Warehouse: 2522 Old Town Road
Union Gap, Washington 98903

