

# Foliar Advantage K

## 2-0-27

### GUARANTEED ANALYSIS

Total Nitrogen (N) .....	2.00%
2.00% Urea Nitrogen	
Soluble Potash (K <sub>2</sub> O) .....	27.00%
Magnesium (Mg) .....	1.00%
Sulfur (S) .....	10.70%


Derived From: Urea, Magnesium Sulfate, and Sulfate of Potash.

**KEEP OUT OF REACH OF CHILDREN**



**WARNING**

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist.



F1548

FOR COMMERCIAL OR PROFESSIONAL USE ONLY

**Loveland**  
PRODUCTS

GUARANTEED BY:

LOVELAND PRODUCTS, INC. • PO BOX 1286 • GREELEY, CO 80632-1286

DENSITY 12.10 LBS/GALLON @ 68° F

NET WEIGHT 30.25 LBS (13.72 kg)

**LIQUID MEASURE 2.5 US GALLONS (9.46 L)**

FoliarAdvantageK\_D0715 product code: XXXXXXXXXX

## GENERAL INFORMATION

**FOLIAR ADVANTAGE K** is a water-soluble foliar nutritional fertilizer of macro and secondary nutrients. It is designed for foliar application and may be applied alone or in tank mixes containing pesticides. However, when use of an unfamiliar mix is made, a compatibility test is always recommended. Foliar fertilization is a supplement to a regular fertilizer program and will not supply the total nutrients required by a crop. **FOLIAR ADVANTAGE K** is recommended as foliar nutrition to be applied to crops where insufficient nutrient levels are indicated by laboratory tissue analysis.

## SHAKE WELL BEFORE USING

### DIRECTIONS FOR USE

Dissolve the desired amount of **FOLIAR ADVANTAGE K** in water before adding to the spray tank. With proper agitation running, add **FOLIAR ADVANTAGE K** to the spray tank. Add **FOLIAR ADVANTAGE K** to the spray tank before adding pesticides. Spray immediately after mixing. Do not store solution mixed with pesticide!

**Alfalfa:** 3-6 pts/acre. Apply after second and fifth cuttings at 4"-6" of re-growth.

**Almonds, Pistachios:** 3-6 pts/acre. 2-3 applications as required from jacket split to hull split as required.

**Apples:** 3-6 pts/acre. 2-3 applications starting at post thinning through early sizing, including 14-21 days post harvest.

**Apricots, Nectarines, Peaches:** 3-4 pts/acre. 2-3 applications from shuck split to full color, including 14-21 days post harvest.

**Avocados:** 3-6 pts/acre. Multiple applications as required. Do not apply within 21 days of harvest.

**Caneberries:** 3-4 pts/acre. 2 applications, one each immediately post bloom and 21-30 days later.

**Celery, Chicory:** 2-4 pts/acre. 2-3 applications during the season as required.

**Cherries, Plums, Prunes:** 3-6 pts/acre. 2-3 applications starting at jacket split to full color, including post harvest.

**Citrus:** 3-6 pts/acre. Multiple applications as required.

**Cotton:** 3-5 pts/acre. 2-3 applications from squaring to full boll development.

**Cucumbers, Melons:** 3-4 pts/acre. 2-3 applications from full bloom through fruit sizing.

**Dry Beans such as Kidneys, Pintos, Limas:** 3-4 pts/acre. 2 applications, one each immediately post bloom and again 30-45 days later.

**Grapes:** 2-4 pts/acre. 2-3 applications from post shatter through early ripening.

**Kiwi:** 3-4 pts/acre. 2-3 applications from early fruit formation to ripening.

### Ornamentals:

**Woody Plants:** 2-4 pts/acre. Multiple applications as required during growing season with one application after September 1st for winter hardiness.

**Pears:** 3-6 pts/acre. 3 applications, one each at early fruit formation, 30 days pre-harvest, and 14-21 days post harvest.

**Peppers, Tomatoes:** 3-5 pts/acre. 2-3 applications from fruit sizing to first color.

**Potatoes:** 3-4 pts/acre. 2-3 applications from flowering through tuberization.

**Strawberries:** 2-4 pts/acre. 2 applications, one each immediately post bloom and 21-30 days later.

**Sugar Beets:** 3-4 pts/acre. 2-3 applications from early beet development at 21-30 day intervals.

**Sweet Potatoes:** 3-4 pts/acre. 2-3 applications from flowering through tuberization.

**Table Grapes:** 3-4 pts/acre. 2-3 applications as required from early fruit formation to ripening.

**Walnuts:** 3-6 pts/acre. 2-3 applications from sizing through packing tissue browning.

For aerial application use a maximum of 3 pints of **FOLIAR ADVANTAGE K** per 10 gallons of water.

Rates suggested are for standard spray volumes of 40-150 gal/acre. For higher volume sprays, rate should be adjusted. **Caution:** Store **FOLIAR ADVANTAGE K** in temperatures between 41°F to 104°F and avoid extreme variations in temperature. A reversible separation of ingredients may occur after long storage, this however does not affect quality or effectiveness of the product.

#### FIRST AID

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

#### STORAGE and DISPOSAL

**STORAGE:** Store in original containers only. Keep containers tightly closed when not in use. Store in a cool, dry well-ventilated area, preferably in a locked storage area away from children, feed and food products, and seed. Store at ambient temperatures. Do not contaminate water, food, or feed by storage or disposal. **DISPOSAL:** Do not reuse container. Plastic: Triple rinse (or equivalent), adding rinse water to spray tank, then offer for recycling at an ACRC site (go to <http://www.acrecycle.org/> for locations) or by reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food, or feed by storage or disposal.

**HANDLING:** Avoid breathing fertilizer mists. Avoid prolonged or repeated skin contact. Avoid eye contact. Wash hands thoroughly after handling this product.

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary. LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when the product is used in accordance with such Directions for Use under normal conditions of use. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that the goods are compatible with all other chemicals under all use conditions.

CALIFORNIA PROPOSITION 65: WARNING: This product contains chemicals known to the State of California to cause cancer or birth defects or other reproductive harm.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lpi.com>

