

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

ForeFront™ R&P

EPA Reg. No. 62719-524

2(ee) Recommendation[†]

For Control of Multiflora Rose

ATTENTION

[†] This recommendation is permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA or state lead pesticide agency.

- It as a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions and limitations on the product label attached to the container for ForeFront™ R&P herbicide.

Use ForeFront™ R&P to control Multiflora rose on rangeland, permanent grass pastures (including grasses grown for hay), and Conservation Reserve Program (CRP) acres.

Specific Use Recommendations

Multiflora rose:

Individual Plant Treatment - Use 42 fl oz (0.3 %v/v) of ForeFront R&P in 100 gal of water with 32 fl oz (0.25% v/v) of a non-ionic surfactant. Or, 32 fl oz (0.25% v/v) of ForeFront R&P can be tank-mixed with Remedy® Ultra at 32 fl oz (0.25% v/v). Apply from full leaf through flowering. For best results, delay treatment for 9-12 months after mowing. Spot treatments may be applied at an equivalent broadcast rate of up to 84 fl oz of ForeFront R&P per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate.

Broadcast - Use 42 fl oz of ForeFront R&P per acre in water with 32 fl oz (0.25% v/v) of a non-ionic surfactant. Or, 32 fl oz of ForeFront R&P can be tank mixed with Remedy Ultra at 16 fl oz. Apply from full leaf through flowering. For best results, delay treatment for 9-12 months after mowing.

™Trademark of Dow AgroSciences LLC

R881-002

Issued: 05/07/07

Initial printing.