

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA REG. NO. 84876-1-279

2(ee) Recommendation
FOR DISTRIBUTION AND USE ONLY IN Florida, Georgia, North Carolina, and South Carolina
IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING
This Product Bulletin Contains New or Additional Directions For Use Which Are Recommended By FMC Corporation Under FIFRA Section 2(ee)

CROP	PEST	RATE OF APPLICATION
Tomato	Southern Blight <i>Sclerotium rolfsii</i>	24.4 – 36.6 fl oz/A (0.4-0.6 lb ai/A)
<p>DIRECTIONS FOR USE: Apply sprays of Fracture® Fungicide following soil applications of Fontelis® Fungicide or other fungicide labeled for Southern Blight control in tomato. Apply Fracture® Fungicide as a directed spray to the soil. Ensure thorough coverage of the lower stems and soil surface surrounding the plants.</p>		
<p>RESTRICTIONS: Do not make more than two sequential applications of Fracture® Fungicide before alternating to a labeled fungicide with a different mode of action. Do not make more than 5 applications of Fracture® Fungicide per harvest cycle.</p>		
<p>PHI: 1 Day</p>		
<p>ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED FRACTURE® FUNGICIDE (EPA REG. NO. 84876-1-279) LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.</p>		



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

FMC and Fracture are trademarks of FMC Corporation.
Fontelis® Fungicide is a trademark of DuPont Crop Protection
©2017 FMC Corporation
Fracture_2ee_Tomato Southern Blight_03-24-2017