

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA

GROUP 21A INSECTICIDE

FujiMite® SC Miticide/Insecticide (EPA Reg. No. 71711-4)

For Use on Stone Fruit (Crop Group 12-12)
States: CA, AZ, WA, OR, HI

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for FujiMite SC Miticide/Insecticide before applying. Use of FujiMite SC Miticide/Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for FujiMite SC Miticide/Insecticide.

FujiMite SC Miticide/Insecticide will provide control of labeled mites when applied by ground at a recommended rate of 2.0 pints per acre.

The following is recommended:

- **Water volume:** Applying by ground using 200 gallons of water per acre or more is highly recommended for full grown trees under heavy mite pressure.
 - If webbing is present, application volume should be increased to ensure that all webbing is penetrated by the spray solution.
 - Control may be reduced if webbing is not adequately penetrated.
- **Adjuvant/spreader:**
 - To optimize control, a 415 oil at a minimum concentration of 1% v/v should be used where a 415 oil is acceptable in a cultural program. If 415 oil is not acceptable, a high-quality silicone surfactant is recommended.
- **Tractor Speed:** For best results, tractor ground speed should not exceed 2 MPH.

©2017 Nichino America, Inc. FujiMite and Nichino America logo are registered trademarks of Nichino America, Inc.

NICHINO
AMERICA[®]
Nichino America, Inc.
4550 Linden Hill Road
Wilmington, DE 19808
888-740-7700

FujiMite SC REC-356-8 07-27-2017
Page 1 of 1