

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA

GROUP 21A INSECTICIDE

FujiMite® SC Miticide/Insecticide (EPA Reg. No. 71711-4)

For Use on Grape
States: CA, AZ, WA, OR, HI

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for FujiMite® SC Miticide/Insecticide before applying. Use of FujiMite SC Miticide/Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for FujiMite SC Miticide/Insecticide.

FujiMite SC Miticide/ Insecticide will provide control of labeled mites when applied by ground at a rate of 2.0 pints per acre.

For Best Results:

- **Water volume:** Apply by ground using a minimum of 50 gallons of water per acre.
 - If webbing is present, application volume should be increased to ensure that all webbing is penetrated by the spray solution.
 - Control may be reduced if webbing is not adequately penetrated.
- **Recommended adjuvant/spreaders:** To optimize control, a silicone-based surfactant should be used. If a silicone based surfactant is not an option due to variety sensitivity, then a quality nonionic surfactant should be used.
- **Tractor speed:** For best results, tractor speed should not exceed 3 MPH. A slower tractor speed of 2 MPH will improve spray coverage and efficacy.

©2020 Nichino America, Inc. FujiMite and Nichino America logo are registered trademarks of Nichino America, Inc.

NICHINO
AMERICA[®]
Nichino America, Inc.
4550 Linden Hill Road
Wilmington, DE 19808
888-740-7700