

## FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA

**GROUP 21A INSECTICIDE**

### FujiMite<sup>®</sup> XLO Miticide/Insecticide (EPA Reg. No. 71711-40)

**For Use on:**

**Small Fruit Vine Climbing Subgroup except Fuzzy Kiwifruit  
(Crop Subgroup 13-07F)  
Berry, Low Growing (Crop Subgroup 13-07G) except Cranberry**

**State: California**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for FujiMite XLO miticide/insecticide before applying. Use of FujiMite XLO miticide/insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for FujiMite XLO miticide/insecticide.

### APPLICATION RATE CHART

<b>Small Fruit Vine Climbing Subgroup except Fuzzy Kiwifruit (Crop Subgroup 13-07F)</b> Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these		
Pest	Rate/ Acre	Use Directions
Mites (see <b>Target Species</b> ) Mealybugs Powdery Mildew (suppression)	2.0 pints	<b>USE RESTRICTIONS</b> <ul style="list-style-type: none"> <li>Apply by ground using a minimum of 50 gallons of water per acre.</li> <li>When using an electro-static sprayer, less than 50 gallons of water per acre may be used; however, do not use less than 5 gallons of water per acre.</li> <li>Do not apply by air.</li> <li>For vines with a heavy canopy, or in high pressure situations, higher water volumes are recommended. If lower water volume amounts are used, tractor speed must be reduced to ensure complete coverage.</li> <li>Do not apply through any type of irrigation system.</li> <li>Do not apply more than 2.0 pints per acre per growing season.</li> <li>Do not make more than 2 applications per growing season.</li> <li>Allow 14 days between applications.</li> <li>Preharvest Interval (PHI): 14 days</li> </ul>
Willamette spider mite	1.5 to 2.0 pints	
Leafhoppers*	1.0 to 2.0 pints	
<small>*Use higher rate for dense foliage. Best control of leafhoppers is achieved by applications when majority of the population is in an immature development stage.</small>		

<b>Berry, Low Growing (Crop Subgroup 13-07G) except Cranberry</b> bearberry; bilberry; blueberry, lowbush; cloudberry; lingonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these		
<b>Pest</b>	<b>Rate/Acre</b>	<b>Use Directions</b>
Mites (see <b>Target Species</b> ) Whiteflies	2.0 pints	<b>USE RESTRICTIONS</b> <ul style="list-style-type: none"> <li>• Apply by ground application using a minimum of 25 gallons of water per acre.</li> <li>• When using an electro-static sprayer, less than 25 gallons of water per acre may be used; however, do not use less than 10 gallons of water per acre.</li> <li>• Do not apply by air.</li> <li>• Do not apply through any type of irrigation system.</li> <li>• Do not apply more than 4.0 pints per acre per crop cycle.</li> <li>• Do not make more than 2 applications per crop cycle.</li> <li>• Allow 14 days between applications.</li> <li>• Preharvest Interval (PHI): 1 day</li> </ul>
<b>NOTE:</b> <ul style="list-style-type: none"> <li>• <b>Temporary pinking of immature green berries may be observed after a FujiMite XLO application on certain strawberry varieties. This effect is transient and does not affect fruit sizing, color or quality.</b></li> <li>• <b>Avoid puddling of spray solution on plastic mulch as this can potentially result in underside scarring of fruit in direct contact with the plastic.</b></li> </ul>		

©2014 Nichino America, Inc. FujiMite and Nichino America logo are registered trademarks of Nichino America, Inc.

**NICHINO**  
**AMERICA**<sup>®</sup>  
 4550 New Linden Hill Road  
 Wilmington, DE 19808  
 888-740-7700