

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA

GROUP 21A INSECTICIDE

FujiMite® XLO Miticide/Insecticide (EPA Reg. No. 71711-40)

For Use on Black Walnut and English Walnut

State: California

Counties: Butte, Colusa, El Dorado, Glenn, Lake, Mendocino, Napa, Placer, Sacramento, Shasta, Siskiyou, Solano, Sonoma, Sutter, Tehama, Trinity, Yolo, Yuba

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for FujiMite® XLO Miticide/Insecticide before applying. Use of FujiMite XLO Miticide/Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for FujiMite XLO Miticide/Insecticide.

FujiMite XLO Miticide/Insecticide will provide control of labeled mites when applied by ground at the rate of 1.5 to 4.0 pints per acre.

For best results:

- Apply when mite populations are beginning to build, before reaching economic thresholds.
- When applying 1.5 pints per acre, 150 gallons of water per acre is recommended.
- When applying 2.0 pints per acre, 200 gallons of water per acre is recommended.
- Thorough spray coverage is necessary; dense foliage or excessive growth will often prevent adequate coverage; adjust spray volumes accordingly.

©2015 Nichino America, Inc. FujiMite and Nichino America logo are registered trademarks of Nichino America, Inc.

NICHINO
AMERICA
4550 New Linden Hill Road
Wilmington, DE 19808
888-740-7700