
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1070A-S3 0917

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

GROUP	1	HERBICIDE
--------------	----------	------------------

Fusilade® DX Herbicide

This supplemental label expires on 08/31/2020 and must not be used or distributed after this date.

Active Ingredient:

Fluazifop-P-butyl

Butyl (*R*)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoate*24.5%

Other Ingredients: 75.5%

Total: 100.0%

Fusilade® DX Herbicide formulation is an emulsifiable concentrate (EC).

*Fusilade DX Herbicide contains 2 pounds (+) isomer (fluazifop-P-butyl) per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1070

All applicable directions, restrictions and precautions on the Fusilade DX Herbicide (EPA Reg. No. 100-1070) label are to be followed. Before using Fusilade DX Herbicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Use of Fusilade DX Herbicide according to this labeling is subject to the limitations imposed by the label affixed to the container for Fusilade DX Herbicide (EPA Reg. No. 100-1070).



DIRECTIONS FOR USE

Fusilade DX Rate Conversion Table for Food Use

Fl oz product/acre	Lb ai fluazifop
6	0.09375
8	0.125
12	0.1875
16	0.250
24	0.375

BLUEBERRY, HIGHBUSH SUBGROUP 13B (As well as Aronia berry; buffalo currant; Chilean guava; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); salal; sea buckthorn; and cultivars, varieties, and/or hybrids of these)

- Do not apply a total of more than 48 fl oz of Fusilade DX Herbicide per acre per year.
- Do not apply more than 24 fl oz of Fusilade DX Herbicide per acre per application.
- Do not apply more than 2 applications at the single maximum application rate per year.
- Maintain a minimum of 14 days between applications.
- Do not harvest within 1 day of last application.

BLUEBERRY, LOWBUSH SUBGROUP 13-07B (As well as lingonberry; currant, native)

- Apply during the non-bearing year or during the year of establishment with the last application no later than 10 months before harvest.
- Do not apply a total of more than 48 fl oz of Fusilade DX Herbicide per acre per year.
- Do not apply more than 24 fl oz of Fusilade DX Herbicide per acre per application.
- Do not apply more than 2 applications at the single maximum application rate per year.
- Maintain a minimum of 14 days between applications.

CANEBERRY SUBGROUP 13-07A (Blackberry; loganberry; raspberry, black and red; wild raspberry; cultivars, varieties, and /or hybrids of these)

- Do not apply a total of more than 48 fl oz of Fusilade DX Herbicide per acre per year.
- Do not apply more than 24 fl oz of Fusilade DX Herbicide per acre per application.
- Do not apply more than 2 applications at the single maximum application rate per year.
- Maintain a minimum of 14 days between applications.
- Do not harvest within 1 day of last application.

LETTUCE, LEAF AND HEAD

- Do not apply a total of more than 48 fl oz of Fusilade DX Herbicide per acre per year.
- Do not apply more than 24 fl oz of Fusilade DX Herbicide per acre per application.
- Do not apply more than 2 applications at the single maximum application rate per year.
- Maintain a minimum of 14 days between applications.
- Do not harvest within 14 days of last application.

ONION, BULB SUBGROUP 3-07A (Daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these)

- Do not apply a total of more than 48 fl oz of Fusilade DX Herbicide per acre per year.
- Do not apply more than 24 fl oz of Fusilade DX Herbicide per acre per application.
- Do not apply more than 2 applications at the single maximum application rate per year.
- Maintain a minimum of 14 days between applications.
- Do not harvest within 45 days of last application.

ONION, GREEN SUBGROUP 3-07B (Chive, fresh leaves; chive, Chinese, fresh leaves; elegans hosta; fritillaria, leaves; kurrat; lady's leek; leek; leek, wild; Onion, Beltsville bunching; onion, fresh; onion, green; onion, macrostem; onion, tree, tops; onion, Welsh, tops; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these)

- Do not apply more than 48 fl oz of Fusilade DX Herbicide per acre per year.
- Do not apply more than 24 fl oz of Fusilade DX Herbicide per acre per application.
- Do not apply more than 2 applications at the single maximum application rate per year.
- Maintain a minimum of 14 days between applications.
- Do not harvest within 14 days of last application.

RHUBARB

- Do not apply a total of more than 32 fl oz of Fusilade DX Herbicide per acre per year.
- Do not apply more than 16 fl oz of Fusilade DX Herbicide per acre per application.
- Do not apply more than 2 applications at the single maximum application rate per year.
- Do not harvest within 14 days of last application.

SMALL FRUIT VINE CLIMBING SUBGROUP 13-07F (EXCEPT FUZZY KIWIFRUIT) (Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these)

- Do not apply a total of more than 72 fl oz of Fusilade DX Herbicide per acre per year.
- Do not apply more than 24 fl oz of Fusilade DX Herbicide per acre per application.
- Do not apply more than 3 applications at the single maximum application rate per year.
- Maintain a minimum of 14 days between applications.
- Do not harvest within 50 days of last application.

STRAWBERRY

- Do not apply a total of more than 16 fl oz of Fusilade DX Herbicide per acre per year.
- Do not apply more than 1 application at the single maximum application rate per year.
- Do not harvest within 14 days of last application.

TUBEROUS AND CORM VEGETABLES SUBGROUP 1D (EXCEPT POTATO) (Arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; sweet potato; taniel; turmeric; yam bean; yam, true)

- Do not apply a total of more than 48 fl oz of Fusilade DX Herbicide per acre per year.
- Do not apply more than 12 fl oz of Fusilade DX Herbicide per acre per application.
- Do not apply more than 4 applications at the single maximum application rate per year.
- Maintain a minimum of 14 days between applications.
- Do not harvest within 14 days of last application.

SPECIAL INSTRUCTIONS FOR USE IN APRICOT; BLUEBERRY, HIGHBUSH (as well as Aronia berry; buffalo currant; Chilean guava; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); lingonberry; salal; sea buckthorn; and cultivars, varieties, and/or hybrids of these); CANEBERRY SUBGROUP (blackberry; loganberry; raspberry, black and red; wild raspberry; cultivars, varieties, and /or hybrids of these); CHERRY, CITRUS FRUITS (calamondin; citrus citron; citrus hybrids; grapefruit; kumquat; lemon; lime; mandarin (tangerine); orange (all); pummelo, and satsuma mandarin); MACADAMIA NUT; NECTARINE; PEACH; PECAN; PLUM; PRUNE; SMALL FRUIT VINE CLIMBING SUBGROUP (EXCEPT FUZZY KIWIFRUIT) (Amur River grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these).

Apply Fusilade DX Herbicide as a directed postemergence application in the interspaces and around the base of plants while avoiding contact of spray with plants. Apply with sufficient spray volume and pressure to ensure complete coverage of target grasses. Applications can be made as a broadcast, strip band, or spot spray application at rates and growth stages listed in Tables 2-5. Refer to the Spray Additives section for recommended spray additives.

Fusilade® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2017 Syngenta