



**FIFRA 24(c) Special Local Need Label (SLN)
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
CALIFORNIA**

For Control of Grasses in Wildlands

FUSILADE® DX Herbicide

EPA REG. NO. 100-1070

EPA SLN No. CA-110010

**This label expires and must not be distributed or used in accordance with this
SLN registration after September 30, 2020**

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA-registered Fusilade® DX Herbicide, EPA Reg No. 100-1070 label.

APPLICATION DIRECTIONS

Apply Fusilade DX Herbicide to wildlands. Make a broadcast application to actively growing targeted grasses following winter rains according to the growth stage and use rate indicated in the **Weeds Controlled** table. Do not apply to grasses which are stressed due to moisture, temperature, low soil fertility, mechanical or chemical injury.

NOTE: Non-target species, particularly grass species, can be injured or killed by Fusilade DX Herbicide applications. It is recommended that a test application be made first to a small representative area to observe any effects on desired species before making applications to larger areas.



Table 1. Weeds Controlled

Grass Species¹	Height (inches)	No. of Leaves Not to Exceed	Rate (Fl. oz/A)
Wild oats (<i>Avena fatua</i>)	2-4	3	24
Perennial ryegrass (<i>Lolium perenne</i>)	2-4	3	24
Ripgut brome (<i>Bromus diandrus</i>)	2-4	3	24
Red brome (<i>Bromus rubens</i>)	2-4	3	24
Soft brome (<i>Bromus hordeaceus</i>)	2-4	3	24

¹ Retreatment at the recommended rate may be needed to control late germinating grasses or if re-growth occurs.

Ground Application – Apply in 20 to 40 gals./A. Use sufficient spray volume and pressure to ensure complete coverage of target grasses.

Spray Adjuvants - Always add one of the following spray adjuvants:

- **Crop oil concentrate** - Add a non-phytotoxic crop oil concentrate or a once refined vegetable oil containing 15-20% approved emulsifier at 0.5-1% v/v (0.5-1 gal./100 gals.) in the finished spray volume.
- **Nonionic Surfactant** - Add a nonionic surfactant containing at least 75% surfactant-active agent at 0.25-0.5% v/v (1-2 qts./100 gals.) in the finished spray volume.

Refer to the EPA-registered label for additional information on application use directions.

GENERAL USE PRECAUTIONS

- Do not apply a total of more than 72 fl. oz. of Fusilade DX Herbicide per acre per year.
- Do not tank mix Fusilade DX Herbicide with triclopyr or aminopyralid containing products as reduced grass control may occur.

RESTRICTIONS:

Chemigation: Do not apply this product to wildlands through any type of irrigation system.

Fusilade® DX Herbicide is a registered trademark of a Syngenta Group Company

©2015 Syngenta

24(c) Registrant
Syngenta Crop Protection, LLC
Greensboro, North Carolina 27419
Phone Number: 1-800-334-9481

EPA SLN No. CA-110010

Label Code: CA1070140BA0915