



# ITERA

## Section 24(c) Special Local Need Label

FLUAZIFOP-P-BUTYL
-------------------

GROUP
-------

1
---

HERBICIDE
-----------

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF HAWAII

**Fusilade® DX Herbicide**  
**For Control of Grasses in Wildlands**

EPA Reg. No. 100-1070  
EPA SLN No. HI-170001

This label expires and must not be distributed or used in accordance with this SLN registration  
after July 11, 2028

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

### APPLICATION DIRECTIONS

Apply Fusilade DX Herbicide to wildlands. Make a broadcast application to actively growing targeted grasses according to the growth stage and use rate indicated in the **Weeds Controlled** table. Do not apply to grasses which are stressed due to moisture, temperature, low soil fertility, mechanical or chemical injury.

**NOTE:** Non-target species, particularly grass species, can be injured or killed by Fusilade DX Herbicide applications. It is recommended that a test application be made first to a small representative area to observe any effects on desired species before making applications to larger areas.

**Table 1. Weeds Controlled**

<b>Grass Species<sup>1</sup></b>	<b>Application Growth Stage</b>	<b>Rate (fl. oz./A)</b>
Bufflegress ( <i>Cenchrus ciliaris</i> )	2-8" tall	24
Southern Sandbur ( <i>Cenchrus echinatus</i> )	2-8" tall	24
Kikuyugrass ( <i>Cenchrus clandestinus</i> )	4-8" runners	24
Finger grass ( <i>Chloris barbata</i> )	2-8" tall	24
Barnyardgrass ( <i>Echinochloa crus-galli</i> )	2-8" tall	24
Goosegrass ( <i>Eleusine indica</i> )	2-8" tall	24
Guineagrass ( <i>Megathyrsus maximus</i> )	2-8" tall	24
Yellow foxtail ( <i>Setaria parviflora</i> )	2-8" tall	24

<sup>1</sup> Retreatment at the recommended rate may be needed to control late germinating grasses or if re-growth occurs.

**Ground and Spot Applications** – Apply Fusilade DX Herbicide at the rate and application growth stage for the listed grass species in Table 1. For broadcast applications, apply in 20 to 40 gallons/A. For Spot Spray applications, refer to the Section 3 Federal label for mixing and application instructions. Use sufficient spray volume and pressure to ensure complete coverage of target grasses.

**Spray Adjuvants** – Always add one of the following spray adjuvants:

- **Crop oil concentrate** – Add a non-phytotoxic crop oil concentrate or a once refined vegetable oil containing 15-20% approved emulsifier at 0.5-1% v/v (0.5-1 gallon/100 gallons) in the finished spray volume.
- **Nonionic Surfactant** – Add a nonionic surfactant containing at least 75% surfactant-active agent at 0.25-0.5% v/v (1-2 qt/100 gallons) in the finished spray volume.

Refer to the Section 3 Federal label for additional information on application use directions.

#### **GENERAL USE PRECAUTIONS**

- Do not apply a total of more than 72 fl oz of Fusilade DX Herbicide per acre per year.
- Do not tank mix Fusilade DX Herbicide with triclopyr or aminopyralid containing products as reduced grass control may occur.

#### **RESTRICTIONS**

- **Chemigation:** Do not apply this product to wildlands through any type of irrigation system.

---

©2023 Syngenta

Fusilade® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, NC 27419-8300

Issue Date: 07/12/2023  
Expiration Date: 07/11/2028

Label Code: HI1070012BA0623