

FIFRA Section 24(c) Special Local Need

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MISSISSIPPI

FOR CONTROL OF GLYPHOSATE-RESISTANT ITALIAN RYEGRASS IN THE FALL
PRIOR TO SOYBEAN OR CORN PLANTING THE FOLLOWING SPRING

GALVAN®

GROUP 5 15 HERBICIDES

EPA REG. NO. 89168-28-89391

EPA SLN NO. MS-220004

ACTIVE INGREDIENTS:	% BY WT.
Metolachlor*	57.64%
Metribuzin**	14.13%
OTHER INGREDIENTS***	28.23%
TOTAL:	100.00%

Contains 5.25 lbs. of metolachlor and 1.25 lbs. of metribuzin per gallon.

*CAS No. 51218-45-2

**CAS No. 21087-64-9

***Contains petroleum distillates

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail).

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the main EPA-registered GALVAN label must be in the possession of the user at the time of application.
- Read the "Conditions of Sale and Limitation of Warranty and Liability" in the label booklet for GALVAN before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

FALL APPLICATION FOR CONTROL OF GLYPHOSATE-RESISTANT ITALIAN RYEGRASS (*Lolium multiflorum*) PRIOR TO SOYBEAN OR CORN PLANTING

Apply GALVAN at 2.5 pints (1.64 lb ai metolachlor and 0.39 lb ai metribuzin) per acre in the fall (September 1 to November 30) after harvest of the previous crop but prior to Italian ryegrass emergence. Regardless of the application date, **DO NOT** apply GALVAN to frozen ground.

A tillage operation may precede the GALVAN application. A fall and/or spring tillage operation may follow application but **DO NOT** exceed an incorporation depth of 2 to 3 inches.

Paraquat at labeled rate can be tank mixed with GALVAN to control emerged Italian ryegrass. Add a nonionic surfactant at 1 quart per 100 gallons (0.25% v/v) in mixtures with paraquat. Read and follow all directions and restrictions on the Paraquat label.

Time Interval Between Treatment with GALVAN and Planting Rotation Crops^{1, 2, 3}

Crop	Months
Alfalfa, winter barley, winter wheat	4.5
Corn, Cotton, peas, rice, spring barley, spring wheat	8
Asparagus, forage grasses, lentils, sainfoin, sugarcane, tomatoes, other crops not listed (except root crops)	12
Onions, sugar beets and other root crops	18
¹ Cover crops for soil building or erosion control may be planted any time, but DO NOT graze or for food or feed. Stand reductions may occur in some areas.	
² Crop rotation recommendations do not include restrictions for the tank mix partner. Refer to the label of the other product for additional restrictions.	
³ Refer to the specific crop use sections for additional crop rotation precautions.	

Restrictions

- **DO NOT** apply more than 2.5 pints (1.64 lb ai metolachlor and 0.39 lb ai metribuzin) per acre per fall application.
- The total amount of this product applied (fall + spring) cannot exceed 3.9 pints (2.56 lb ai metolachlor and 0.61 lb ai metribuzin) pints per acre per year.
- Make no more than 1 fall application of this product.

24(c) Registrant:

Innvictis Crop Care, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

All trademarks are the property of their respective owners.

022723