

Specimen Label



Insecticidal Bait

*Trademark of Dow AgroSciences LLC

For selective attractance and control of multiple species of tephritid fruit flies infesting various tree, fruit, nut, vine and vegetable crops and ornamentals, and on non-crop vegetation which may serve as resting sites

Active Ingredients:

spinosad (a mixture of spinosyn A and spinosyn D)	0.02%
Inert ingredients: includes water, sugars and attractants	99.98%
Total	100.00%

Ready to use formulation.

U.S. Patent No. 5,362,634 and 5,496,931

EPA Reg. No. 62719-359

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Swallowed

Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or when disposing of equipment washwaters.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

AVOID FREEZING

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: There are no specific entry requirements following application of this product when applied by government agencies in area-wide eradication and prevention programs.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site according to label use directions or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

GF-120 Naturalyte* Fruit Fly Bait attracts and controls multiple species of tephritid fruit flies infesting various tree, fruit, nut, vine and vegetable crops and ornamentals, and on non-crop vegetation which may serve as resting sites. Aerial or ground applications in production agriculture or directed ground applications to individual plants are permitted. In urban areas, however, use is limited to directed ground applications.

Chemigation: Do not apply through any type of irrigation equipment.

GF-120 may be used on the following crops, ornamentals and on non-crop vegetation which may serve as resting sites for adult flies:

Bushberry including, but not limited to: blueberry, currant, elderberry, gooseberry, huckleberry, junberry, salal, and lingonberry

Citrus including, but not limited to: oranges, grapefruit, lemons, limes, kumquat, and tangerines

Cole Crops including, but not limited to: broccoli, Brussels sprouts, cabbage, Chinese cabbage (bok choy, napa, etc.), cauliflower, Cavalo, collards, kale, kohlrabi, mustard greens, mustard spinach, Chinese mustard cabbage (gai choy), and rape greens

Corn (field, sweet and seed)

Cucurbit Crops including, but not limited to: cucumber, squash, muskmelons (cantaloupe, honeydew, etc.) pumpkin, and watermelons

Fruiting Vegetable Crops including, but not limited to: eggplant, pepper, and tomato

Leafy Vegetables including, but not limited to: head and leaf lettuce, celery, arugula, chervil, edible chrysanthemum, cilantro, cress, dandelion, endive, radicchio, rhubarb, spinach, and turnip greens

Ornamentals and Non-Crop Foliage: along field edges and adjacent to treated areas that may serve as a resting sites for adult flies.

Pome Fruit including, but not limited to: apples, pears, crabapple, loquat, quince, and mayhaw

Potatoes and Tuberous and Corm Vegetables including, but not limited to: cassava, chayote root, Chinese artichoke, Jerusalem artichoke, potatoes, sweet potatoes, and yams

Stone Fruit including, but not limited to: peaches, plums, cherries, nectarines, prunes, and apricots

Succulent and Dried Beans and Peas including, but not limited to: adzuki bean, fava bean, garbanzo bean, field bean, kidney bean, snap bean, wax bean, blackeyed pea, crowder pea, English pea, garden pea, snow pea, and sugar snap pea

Tree Nuts including, but not limited to: walnuts and pistachios

Tropical Fruit including, but not limited to: sugar apple, cherimoya, atemoya, custard apple, ilama, soursop, biriba, lychee, longan, Spanish lime, rambutan, pulasan, papaya, star apple, black sapote, mango, sapodilla, canistel, mamey sapote, avocado, guava (kelly), feijoa, jaboticaba, wax jambu, starfruit, passionfruit, acerola, and white sapote.

Preparation for Use

Mixing - GF-120 Naturalyte Fruit Fly Bait is a bait concentrate that should be diluted with water. The most effective dilution for aerial and most other applications is a 1:1.5 (GF-120:water) dilution (e.g., to make 10 liters of spray solution, add 6 liters of water to 4 liters of GF-120). For ground applications and applications in low relative humidity, dilutions of up to 1:5 (GF-120:water) can be made. First add water (one-half of the volume to be mixed) to the spray tank or premixing tank and start the agitation system. Then add the full amount of GF-120 followed by an equal amount of water. If a full container of GF-120 is used, the empty container should then be triple rinsed by filling it one-third full with water, shaking well and adding the rinsate to the spray tank. Repeat two more times so the container is triple rinsed and then complete filling the spray tank until the proper dilution is obtained. Constant agitation of the spray solution is recommended to ensure uniformity of spray mixture. Allow agitation system to operate for at least 5 minutes before applying. Once diluted, GF-120 should be used within 24 hours. Concentrated GF-120 will not settle and does not need to be shaken before mixing.

Application

Proper application techniques help ensure adequate coverage and correct dosage necessary to obtain optimum control of insect pests. A large spray droplet size of 4000 to 6000 μ (4-6 mm) is recommended to optimize length of bait attractance. Fruit flies can detect the bait from several yards away. When aerially applying use ULV applications but with coarse nozzles that will produce the desired droplet size and target 20 to 80 droplets per square meter. By ground, spot or strip spray several areas on the inner canopy of fruiting plants. Avoid weather conditions that could result in drift to nontarget areas. Direct spray application to bottoms of leaves and leaves inside the foliage canopy to reduce direct exposure to sun and rain. This product resists wash off, but will lose effectiveness if exposed to rain and overhead irrigation. When possible, potential for rain or irrigation schedules should be considered when planning applications. Begin applications as soon as monitoring traps indicate flies are present or 2 to 3 weeks before fruit begins to ripen. Repeat applications every 7 to 14 days, shortening the application interval during rainy periods and as fruit ripens. Remove fruit as soon as ripe, particularly any overly ripe fruit on the tree or ground.

Pests and Application Rates:

Pests Controlled or Suppressed	Amount of Undiluted GF-120 Spray Solution ¹			
	Broadcast Application		Spot Spray of Individual Plants	
	fl oz/ acre	liters/ hectare	fl oz/ tree	ml/ tree
tephritid fruit flies (including but not limited to:) apple maggot Caribbean fruit fly cherry fruit fly Mediterranean fruit fly melon fly Mexican fruit fly olive fly Oriental fruit fly walnut husk fly	10 - 20	2 - 4	1 - 3	30 - 90

¹ Choose rate based on pest pressure and amount of foliage needed to cover.

Amount of Final Spray Solution for Different Rates and Dilution Ratios:

Dilution Ratio GF-120:water	Rates of GF-120 per Acre				
	10 fl oz/acre	12 fl oz/acre	15 fl oz/acre	18 fl oz/acre	20 fl oz/acre
1:1.5	25	30	37.5	45	50
1:2	30	36	45	54	60
1:3	40	48	60	72	80
1:4	50	60	75	90	100
1:5	60	72	90	108	120

*Entries in the table are the amount of final diluted spray solution per acre

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Label Code: D02-134-002

Replaces Label: D02-134-001

LOES Number: 010-01760

'[

EPA Accepted: 11/13/02

Revisions:

- 1. Added the following crops: bushberries, kumquat (to citrus), pome fruit, tree nuts.