

PRODUCT BULLETIN
2(ee) Recommendation

THIS PRODUCT BULLETIN CONTAINS NEW OR
ADDITIONAL DIRECTIONS FOR USE
WHICH ARE RECOMMENDED BY INNVICTIS
CROP CARE, LLC UNDER FIFRA SECTION 2(ee)

GOVEE™

EPA Reg. No. 89167-90-89391

2(ee) Recommendation†

**For Weed Control in Fallow Beds to be Planted to Soybeans, Corn, or Cotton in AL, AR, FL, GA,
LA, MO, MS, NC, OK, SC, TN, VA**

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions and limitations on the product label attached to the container for GOVEE.

DIRECTIONS FOR USE

GOVEE, alone or in tank mix combination with glyphosate, paraquat, glufosinate, 2,4-D or dicamba is effective for the control of winter annual broadleaf weeds in fallow beds to be planted to soybeans, corn or cotton. Once a fallow bed application is made, the soil surface should be left undisturbed for the period of time in which weed control is desired. Cultural practices that disturb or redistribute surface soil following treatment with GOVEE will reduce weed control effectiveness. If a fallow bed treatment is applied thirty or more days before planting soybeans, corn or cotton and at least three significant rainfalls (0.25 inch or greater) have occurred following application, then soybean, corn or cotton can be planted directly into the stale seedbed without incorporation. If three significant rainfalls (0.25 inches or greater) have not occurred during the 30 or more days before planting, then GOVEE must be incorporated into the soil to a depth of 2.5 inches or more prior to planting soybean, corn or cotton.

Exercise extreme care to avoid herbicide contact with any desirable dormant or non-dormant crop, plant, tree or vegetation as severe injury may result.

Weeds Controlled: GOVEE provides preemergence and postemergence control of the following weeds when used at recommended dosages and weed stage. †

Amaranth, Palmer
Geranium, Carolina
Henbit

Ladysthumb
Mustard Species
Pigweed, Redroot

Sida, Prickly
Smartweed
Waterhemp

† Thorough spray coverage is essential for optimum post-emergence activity of GOVEE. For optimum postemergence control, a tank mixture of GOVEE with glyphosate, 2,4-D, glufosinate, dicamba, or paraquat is recommended.

Application Rates (GOVEE Used Alone)

GOVEE may be applied at 1 to 2 pints (0.25 to 0.5 lb ai) per broadcast acre. The lower rate (1 pint per acre) should provide up to 4 weeks of preemergence control of susceptible weeds and provide postemergence control of susceptible weeds (up to 4-leaf stage). The higher rate (2 pints per acre) should provide preemergence control of susceptible weeds for up to 8 weeks and postemergence control of susceptible

weeds (up to 6-leaf stage). The best preemergence control is achieved when irrigation or rainfall occurs within 3 or 4 weeks following application.

Tank Mixes with GOVEE

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Application Rates (GOVEE Used in Combination with Other Products): GOVEE can be tank mixed with glyphosate, glufosinate, paraquat, 2,4-D or dicamba to obtain postemergence control of annual grass weeds, volunteer grains and broadleaf weeds. One to 2 pints (0.25 to 0.5 lb ai) per acre of GOVEE may be tank mixed with labeled rates of glyphosate, glufosinate, paraquat, 2,4-D or dicamba. Apply at the recommended rates and growth stages for susceptible weed species and in a manner consistent with the tank mix product label.

For enhanced contact activity (burndown/suppression) with glyphosate, glufosinate, paraquat, 2,4-D or dicamba, GOVEE at the rate of 3 to 7 fluid ounces (0.05 to 0.1 lb ai) per acre may be added to labeled rates of glyphosate, glufosinate, paraquat, 2,4-D or dicamba. Apply at the recommended rates and growth stages for susceptible weed species and in a manner consistent with the tank mix product label.

Method of Application (Ground or Aerial Application)

Apply GOVEE in a spray volume of 15 or more gallons per acre with ground equipment. Increase the spray volume, if necessary, to ensure thorough coverage of existing weeds. Use a spray volume of 10 or more gallons per acre (5 gallons or more for glyphosate tank mix) when aerially applying. Carefully follow aerial application requirements and advisory information in the Fallow Bed section of the product label for GOVEE.

Fallow Bed - Specific Use Restrictions

In addition to the following, also observe use restrictions listed at the beginning of the GOVEE label.

- **DO NOT** apply more than 2 pints (0.5 lb ai) of GOVEE per acre per fallow season.
- **Chemigation:** Under this 2(ee) label, **DO NOT** apply this product through any type of irrigation system.
- GOVEE can be applied at 1 to 2 pints (0.25 to 0.5 lb ai) per acre as a fallow bed application at 30 days or more prior to planting soybeans, corn or cotton with the following requirements:
 - If three activating rainfalls or irrigation (0.25 inches or greater) **HAVE** been received after application during the period of 30 days or greater between application and soybean, corn or cotton planting, then the crop can be planted directly into the stale seedbed without GOVEE incorporation.
 - If three activating rainfalls or irrigation (0.25 inches or greater) have **NOT** been received after application during the period of 30 days or greater between application and soybean, corn or cotton planting, then GOVEE must be incorporated into the soil to a depth of 2.5 inches or more prior to planting soybean, corn or cotton.
- **DO NOT** use corn plants from a treated field for green chop, ensilage, forage or fodder.
- GOVEE is phytotoxic to plant foliage. **DO NOT** apply when weather conditions favor drift. Avoid drift to all non-target areas.
- The use of any plants from a treated field for green chop, ensilage, forage or fodder or the feeding or grazing of animals on any treated area is prohibited.

Distributed By:

INNICTIS CROP CARE, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

100121

All trademarks are the property of their respective owners