



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF GRANDEVO® CG BIOINSECTICIDE
ON TROPICAL AND SUBTROPICAL FRUIT FOR MANAGEMENT OF MITES AND MEALYBUGS

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(Ee) RECOMMENDATIONS ARE RECOGNIZED.

Grandevo® CG
EPA Reg. No. 84059-27

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Target Arthropods	Application Method	Product Use Rate per Application	Application Instructions
Tropical and Subtropical Fruit	<i>Tetranychid</i> sp. and <i>Oligonychid</i> sp. mites <i>Pseudococcus</i> sp. mealybugs	Foliar	2 lb-3 lb/Acre	Begin applications at first incidence of pest presence and continue on a 5 to 10-day schedule until harvest or when pest activity is no longer apparent. Apply at sufficient water volume to ensure thorough coverage and apply a surfactant at labelled rate.
GRANDEVO® CG has a pre-harvest interval (PHI) of 0 days. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.				

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

Grandevo® CG is a registered trademark of Marrone Bio Innovations, Inc.
© Copyright Marrone Bio Innovations, Inc., 2022
Marrone Bio Innovations, Inc.
1540 Drew Ave.
Davis, CA 95618
Telephone (530) 750-2800
www.marronebioinnovations.com