

SUPPLEMENTAL LABEL

FLUTIANIL | GROUP | U13 | FUNGICIDE



EPA Reg. No. 11581-6-71711

For Use On: Melon Subgroup 9A and Squash/Cucumber Subgroup 9B

Not for Sale, Sale Into, Distribution and/or Use in Nassau and Suffolk Counties of New York State

This supplemental label expires January 17, 2023 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for GATTEN® Fungicide before applying. Use of GATTEN Fungicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for GATTEN Fungicide.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

DIRECTIONS FOR APPLICATION

Melon Subgroup 9A and Squash/Cucumber Subgroup 9B

chayote (fruit); Chinese waxgourd (Chinese preserving melon); cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); muskmelon (hybrids and/or cultivars of *Cucumis melo*) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); pumpkin; squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

Target Disease	Application Rate fl oz product/A (lb ai/A)	PHI	Directions
<i>G. cichoracearum</i> (Powdery Mildew)	6.0 (0.0198) to 8.0 (0.0264)	0	<ul style="list-style-type: none">• Use water volumes of 30 to 100 gallons per acre for sprays.• Repeat applications at 7 to 14-day intervals, depending on disease pressure.• Apply to foliage during the seedling to the fruiting stage using broadcast ground equipment.

Restrictions

- **Maximum Application Rate:** 8.0 fl oz product/A (0.0264 lb ai/A)
- **DO NOT** apply more than 5 applications during each growing season.
- Maximum of two growing seasons per year.
- **DO NOT** exceed a maximum of 0.132 lb ai per acre per season.
- Crop can be harvested after product has dried.
- **NEW YORK: DO NOT** apply more than 16 fl oz product/acre per year. **DO NOT** exceed 0.0528 lb ai/acre per year.

©2020 OAT Agrio Co. LTD. Gatten is a registered trademark of OAT Agrio Co., Ltd. Nichino America logo is a registered trademark of Nichino America, Inc.

NICHINO
AMERICA

Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700