

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Goal[®] 2XL herbicide

EPA Reg. No. 62719-424

Special 2(ee) Recommendation†

Fallow-Bed Use Prior to Transplanting Strawberries or Peppers Grown in Plastic Culture

(For distribution and use only in the state California)

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been reviewed or approved by EPA.

- This recommendation must be in the possession of the user at the time of application.
- Read the label affixed to the container for Goal[®] 2XL herbicide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Except as described in this product bulletin, use of Goal 2XL is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Goal[®] 2XL herbicide may be applied broadcast or banded as a fallow bed application to pre-formed beds prior to planting of strawberries or peppers grown in plastic culture.

It is recommended that soil moisture be used to activate Goal 2XL using one of the following practices soon after application.

- Irrigate the beds with ½ inch of sprinkler irrigation and then put plastic down anytime during the 30-day treatment-to-planting interval.
(or)
- If there is adequate existing soil moisture, apply plastic to the beds as soon as possible after application and allow the moisture which condenses and accumulates beneath the plastic to thoroughly wet the treated soil.

Mechanical incorporation of the fallow-bed treatment prior to laying plastic is not required. Not disturbing the soil surface may allow for extended weed control. Not incorporating increases the potential for crop injury, especially under wet conditions. Therefore, the treatment should be incorporated if the risk of crop injury is not acceptable. Follow the minimum treatment-to-planting intervals outlined below (also found on the main product label).

Minimum Treatment-to-Planting Interval:

Transplanted Crops	Goal 2XL Use Rate	
	Up to 1 pt/acre	Up to 2 pt/acre
Pepper	30 days	30 days
Strawberries	30 days	30 days

®Trademark of Dow AgroSciences LLC

R-843-003
Issued: 12-02-2004
Initial printing.