

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Goal* 2XL

Dow AgroSciences EPA Reg. No. 62719-424

24(c) Special Local Need Registration SLN MS-020016

For Use in Deciduous Tree Plantings
(For Distribution and Use Only in the State of Mississippi)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Goal* 2XL herbicide before applying. Follow all other applicable directions, restrictions, and precautions on the EPA-registered label.
- Use of Goal 2XL according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Goal 2XL.
- Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions, and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

Directions for Use

Goal 2XL herbicide is an effective herbicide for preemergence and postemergence control of certain broadleaf weeds in deciduous tree plantings. In new plantings, over-the-top applications of Goal 2XL herbicide should be made soon after transplanting of dormant deciduous tree seedlings. Subsequent applications can be made to nondormant, fully foliated trees. Over-the-top applications to nondormant trees may result in minor discoloration and spotting of the foliage; however, trees will outgrow this condition. Leaves that are fully expanded and hardened off will exhibit less injury than newly emerged leaves or new bud growth. Applications to newly emerged leaves and/or new bud growth will result in leaf injury and is done at the user's risk.

Dosage

Goal 2XL herbicide may be applied at 2 to 8 pints (0.5 to 2lb active) per broadcast acre for preemergence and postemergence weed control. A maximum of 8 pints of Goal 2XL herbicide (2.0 lb active) per acre per season may be applied as a result of single or multiple applications. The addition of 1 quart of LA TRG~ A G-98 or a comparable 80% active nonionic surfactant per 100 gallons of spray mix will assist in spray coverage and wetting of weeds for postemergence coverage.

Deciduous Species

Cottonwood, Eastern	Populus deltoides
Oak, Cherrybark	Quercus pagoda
Oak, Nutt All	Quercus nuttallii
Sweetgum	Liquidambar styracifula
Sycamore	Platanus occidentalis

(Selected deciduous trees listed on this label have shown tolerance to Goal 2XL herbicide. It is impossible, however, to evaluate this product on all varieties, biotypes and cultivars of listed species on this label under all possible growing conditions. The user should exercise reasonable judgment and caution with this product. Until familiar with results under use growing conditions, limit application of this

product to a small treated area to determine plant tolerance and extent of injury if such occurs, prior to initiating large-scale applications.

Weeds Controlled

When Goal 2XL herbicide is applied preemergence or postemergence to weed seedlings (not exceeding 4- to 6-leaf stage) at recommended dosages, the following broadleaf weeds are controlled:

Croton, Tropic	Nightshade, Black
Groundcherry, Cutleaf	Nightshade, Hairy
Groundcherry, Wright	Pigweed, Redroot
Groundsel, Common	Purslane, Common
Jimsonweed	Sesbania, Hemp
Knowtweed, Prostrate	**Sida, Prickly (Tea Weed)
Morningglory, Annual	Smartweed, Pennsylvania
Mustard, Hedge	Velvetleaf
Nightshade, American Black	

*Postemergence up to 4- to 6-leaf stage.

**Highest rate or multiple applications may be required for acceptable control.

Timing and Method of Application

For optimum weed control, a dormant over-the-top application of Goal 2XL herbicide should be made prior to weed seedling emergence followed by a postemergence application after tree foliage have fully expanded. For weed management programs using only a single application per season, a preemergence application of Goal 2XL herbicide is recommended.

Goal 2XL herbicide should be applied at 20 to 40 psi in a minimum of 20 gallons of water per acre depending upon density of emerged weeds. Spray volume should be increased as weed height and density increase. Use a low-pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

Mixing Directions

Fill the spray tank at least one-third full of clean water. With the pump and agitator running, add the recommended amount of herbicide to the spray tank. Complete filling of the spray tank with water. Maintain agitation until spraying is complete.

Chemigation: Do not apply this product through any type of irrigation system.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

This product is highly toxic to aquatic invertebrates, aquatic plants, wildlife and fish. Use with care when applying in areas frequented by wildlife or adjacent to any body of water or wetland area. Do not apply when weather conditions favor drift or erosion from target areas.

24(c) Registrant: Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

*Trademark of Dow AgroSciences LLC

Revisions: Section 24(c) for 2002.

D06-843-043
Approved: 07/17/2002
Replaces D06-843-037