

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Goal[®] 2XL

Dow AgroSciences EPA Reg. No. 62719-424

24(c) Special Local Need Registration SLN HI-060002

For Weed Control in First Year Sugarcane Only

(For Distribution and Use Only in the State of Hawaii)

Conditions and Risks of Use for Special Local Need

USE OF Goal[®] 2XL HERBICIDE (THE "PRODUCT") ON SUGARCANE (THE "CROP") FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP.

This Product is available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Supplemental Labeling, you acknowledge and accept that to the extent permitted by law:

- (1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Dow AgroSciences does not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental labeling;
- (3) Dow AgroSciences does not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- (4) Dow AgroSciences disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) these Conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Dow AgroSciences, or its respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

These Conditions and Risks of Use for Special Local Need are required by Dow AgroSciences and not specified by the US EPA or the State of Hawaii.

Expiration Date: Valid until November 15, 2011 or until otherwise amended, withdrawn, cancelled or suspended.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Goal[®] 2XL herbicide before applying. Follow all other applicable directions, restrictions, and precautions on the EPA-registered label.
- Use of Goal 2XL according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Goal 2XL.
- Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions, and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

Directions for Use

General Information

Goal 2XL provides preemergence control of many broadleaf and grass weeds during the establishment of sugarcane, including Aiea morningglory, guineagrass and triazine-resistant swollen fingergrass.

Rates and Method of Application

Apply 1 to 8 pints per acre (0.25 to 2.0 lbs. active) as a broadcast spray before sugarcane shoot leaves unfurl. The lowest rate can be used prior to weed emergence when the soil is moist. Most drip-irrigated fields will require 2-4 pints per acre (0.5 to 1.0 lbs. active). Use higher rates for preemergence control in dry weather and to control emerged weeds.

Apply 1 to 8 pints per acre (0.25 to 2.0 lbs. active) as a directed spray after the sugarcane leaves have unfurled. Avoid contact with the leaves of the sugarcane or crop damage will result.

Goal 2XL can also be used in combination with glyphosate as a directed spray field edge application if emerged weeds are present.

Goal 2XL herbicide should be thoroughly mixed with clean water at recommended concentration and applied using a low-pressure ground sprayer equipped in a minimum spray volume of 20 gallons per acre.

If spray drift or treated soil come into contact with emerging sugarcane spikes or whorls, crop injury may occur. The crop will outgrow this temporary injury, and the effect generally is offset by the benefit of weed control.

Weeds Controlled

Goal 2XL herbicide will provide preemergence control/suppression of the following weeds when applied at the recommended dosage:

Aiea morningglory
guineagrass
swollen fingergrass

Crop-Specific Use Restrictions

The following use restrictions must be observed when Goal 2XL herbicide is used as recommended on this label:

- Follow the General Use Restrictions on the EPA-registered label.
- For wind speeds at or below 10 mph, see package label for spray drift precautions.

- For wind speeds between 10-20 mph, **do not apply this product in a manner that allows spray to drift from the application target site and cause harm to humans, animals or other non-target sites.** Recommended applications to reduce off-site drift include, but are not limited to, the following:
 - Apply by ground boom with nozzle height no more than 2 feet above ground for broadcast applications or no more than 1.25 foot above the ground for directed sprays,
 - Use coarse spray according to ASAE 572 definition for standard nozzles or VMD of 475 microns for spinning atomizer nozzles.
 - Use hooded or shielded sprayers or other means to reduce drift.
 - Use spray drift retardants and/or other measures known to control drift.
- This labeling must be in the possession of the user at the time of application.
- For use on non-bearing sugarcane only.
- Do not apply more than 8 quarts Goal 2XL per acre (4 lbs. active) per crop as result of multiple applications.
- **Preharvest Interval:** Do not harvest for food or feed within one year of application.
- Do not apply more than a total of 4.0 lbs of oxyfluorfen active ingredient per acre per year.
- Avoid contact with the leaves of the sugarcane or crop damage will result.
- **Chemigation:** Do not apply this product through any type of irrigation system for uses described in this supplemental labeling.
- **Do not harvest for food or feed within one year of application.**

Issue Date: November 16, 2006
Revision Date: May 19, 2010

®Trademark of Dow AgroSciences LLC

R843-100
Approved: 06/14/10
Replaces: R843-073