

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS– NOT TO BE
USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A
CERTIFIED APPLICATOR.**

syngenta.

INTIRA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TENNESSEE

**Gramoxone® SL 3.0
For Weed Control in Row Middles of Lima Bean and Southern Pea**

**EPA Reg. No. 100-1652
EPA SLN No. TN-210006**

**This label expires and must not be distributed or used in accordance with this SLN
registration after June 2, 2026**

**KEEP OUT OF REACH OF CHILDREN /
MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.**



**DANGER / POISON
PELIGRO / VENENO**

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

Target Weed	Rate pt/A	Application Timing	Use Directions
Palmer amaranth, other annual grass and broadleaf weeds	1.3-2.0	Postemergence (Directed with Hooded sprayer)	<p>Apply as a post-directed spray between the rows using a hooded sprayer. Apply when weeds are 2-4 inches tall and actively growing, larger weeds may not be adequately controlled. Use the higher rate on larger or hard to control weeds. Apply in a minimum of 10 gal/A of water.</p> <p>Use nozzles that provide uniform coverage within the treated area. Keep hoods on these sprayers adjusted to protect desirable vegetation. Operate equipment in a manner that avoids bouncing or raising the hoods off the ground. Tractor operating speed must be adjusted to avoid bouncing of the spray hoods.</p> <p>Add either: 1) a nonionic surfactant containing 80% or more surface-active agent at 0.25% v/v (2 pt/100 gal) of the finished spray volume OR 2) Crop Oil Concentrate or Methylated Seed Oil containing 15-20% approved emulsifier at 1.0% v/v (1 gal/100 gal) of the finished spray volume.</p> <p>A second application may be made, allow a minimum of 7 days between applications.</p>
Precaution: <ul style="list-style-type: none"> Spray should not contact green stems, foliage, or pods, as injury may result. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Maximum Single Application Rate: 2.0 pt/A REI: 24 hours Minimum Application Interval: 7 days Maximum Annual Rate: 4.0 pt/A DO NOT make more than 2 applications per year Pre-Harvest Interval (PHI): 21 days 			

©2021 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company.

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: TN1652034AA0521