

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY**

**FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS ONLY– NOT TO BE
USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A
CERTIFIED APPLICATOR.**

syngenta.

ENTRA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS

**Gramoxone® SL 2.0
Harvest Aid Applications on Cotton in Texas
(Except in the Counties of Starr, Hidalgo, Willacy, and Cameron)**

**EPA Reg. No. 100-1431
EPA SLN No. TX-220002**

**This label expires and must not be distributed or used in accordance with this SLN
registration after December 31, 2023**

**KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS
NIÑOS.**



**DANGER / POISON
PELIGRO / VENENO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

Specific Crop Use Directions

Three multiple broadcast harvest aid applications of Gramoxone SL 2.0 may be used. **Do not exceed a total of 3 pints per acre per crop season.** Use a minimum total spray volume per acre of 10 gallons by ground, and 5 gallons by air.

| Crop | Rate per Acre | Precautions, Restrictions and Comments |
|--|--|---|
| Tank mixes with phosphate or chlorate defoliant | 8.1 to 16.1 fl oz alone <u>or</u> tank mixed with 1 1/2 to 2 gal chlorate, <u>or</u> 1 1/2 to 2 pt phosphate | <ul style="list-style-type: none"> Do not harvest within 7 days of application when used with phosphate or chlorate defoliant. In desert cotton areas or on rank vigorous cotton, use the 16.1 fl oz rate of Gramoxone SL 2.0 Apply when 75% or more of the bolls are open and remaining bolls to be harvested are mature. Development of immature bolls will be inhibited. Follow Harvest Aid Use Precautions listed on the Federal label. |
| Tank mixes for boll opening and defoliation of southern cotton | 4.1 to 5.0 fl oz | <ul style="list-style-type: none"> To aid in defoliation and opening of mature bolls, Gramoxone SL 2.0 may be tank mixed with the following products: DEF[®] defoliant, DROPP[®] defoliant, PREP[™] PGR, FOLEX[®] defoliant. Apply when 60% or more of the bolls are open and the remaining bolls to be harvested are mature. Development of immature bolls will be inhibited. Refer to tank mix product label for rate, directions, limitations and caution. |
| Harvest aid for defoliation and boll opening of stripper or spindle harvested cotton | 3.1 to 11.2 fl oz | <ul style="list-style-type: none"> Apply when 75% of the bolls are open and the remaining bolls to be harvested are mature. Development of immature bolls will be inhibited. Slice bolls and inspect the seed for maturity. Gramoxone SL 2.0 may be applied alone or tank mixed with the cotton harvest aid products listed above. To avoid leaf sticking, apply Gramoxone SL 2.0 as a desiccant approximately 3-7 days after defoliation or a conditioning application and 7-14 days before harvest. Cooler temperatures may cause a longer waiting period between application of Gramoxone SL 2.0 as a desiccant and defoliation/conditioning. Lower rates in the range may be necessary south of I-10 in Texas where temperatures are typically higher during defoliation. |
| Harvest aid desiccation of stripper cotton or spindle harvested cotton late season | 16.8 to 31.5 fl oz | <ul style="list-style-type: none"> Use to desiccate regrowth occurring after defoliation or desiccation. Apply when 85% of the bolls are open and the remaining bolls to be harvested are mature. Development of immature bolls will be inhibited. Slice bolls and inspect the seed for maturity. To avoid leaf sticking, apply Gramoxone SL 2.0 |

| Crop | Rate per Acre | Precautions, Restrictions and Comments |
|-------------------------|--------------------|--|
| | | as a desiccant approximately 3-7 days after defoliation or a conditioning application and 7-14 days before harvest. <ul style="list-style-type: none"> Lower rates in the range may be necessary south of I-10 in Texas where temperatures are typically higher during defoliation. |
| Desiccation of Regrowth | 16.8 to 31.5 fl oz | <ul style="list-style-type: none"> Use to desiccate regrowth after defoliation or desiccation. Regrowth is difficult to control, therefore, thorough coverage with the full recommended rate is necessary. Control is dependent upon growing condition and desiccation of small new growth may not always be complete. Use higher rate if regrowth is excessive. |

Refer to the Gramoxone SL 2.0 federal label for additional cotton recommended uses and nonionic surfactant recommendations.

Additional Precautions, Restrictions and Comments:

- It is advisable, because of extremes in environmental and plant conditions, to apply the range of rates on a small block of cotton to determine the rate that best fits your needs.
- Do not pasture lactating dairy animals.
- Do not pasture livestock in treated fields or feed treated foliage.
- When combined with phosphate defoliant, observe livestock cautions listed on the phosphate label.
- Do not harvest within 7 days of application when used with phosphate or chlorate defoliant.
- Do not apply this product through any type of irrigation system.

©2021 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company.
 Def®, Dropp®, Prep™ PGR trademarks of Bayer CropScience.
 Folex® trademark of Amvac Chemical Corporation.

24(c) Registrant:
 Syngenta Crop Protection, LLC
 P. O. Box 18300
 Greensboro, NC 27419-8300

Label Code: TX1431107CA0921