

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

syngenta.

TIARA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MISSISSIPPI

**Gramoxone SL 2.0
For Preplant / Preemergence Control of Weeds in Peanuts**

**EPA Reg. No. 100-1431
EPA SLN No. MS-190002**

Active Ingredient:

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride)30.1%

Other Ingredients: 69.9%

Total: 100.0%

**This label expires and must not be distributed or used in accordance with this
SLN registration after December 31, 2024**

KEEP OUT OF REACH OF CHILDREN



**DANGER / POISON
PELIGRO / VENENO**

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
- Failure to follow the Directions for Use and Precautions on the EPA-registered Gramoxone SL 2.0 label may result in poor pest control, crop injury, or illegal residues.

PEANUTS		
Rate pt/A	Application Timing	Use Directions
1.9 – 3.75	Preplant / Preemergence	<p>Apply to burndown weeds in peanut fields preplant / preemergence.</p> <p>Apply as a ground broadcast application in a minimum of 10 gal/A of spray volume. Increase carrier volume for dense weed populations and larger weeds</p> <p>Use the higher rate on dense populations and/or on larger or harder to control weeds. Apply when weeds are actively growing and between 1" to 6" in height. Vegetation 6" or taller may not be controlled.</p> <p>Always add either: 1) Nonionic surfactant containing 80% or more surface agent at 0.25% v/v (2 pints per 100 gallons), of the finished spray volume, OR 2) crop oil concentrate (non phytotoxic) containing 15 to 20% approved emulsifier, at 1% v/v (1 gallon per 100 gallons) of the finished spray volume.</p> <p>Tank Mix Options: Gramoxone SL 2.0 may be tank-mixed with other herbicides registered for this use for improved weed control.</p>
<p>Use Restrictions:</p> <ol style="list-style-type: none"> Maximum Single Application Rate: 3.75 pt/A. Maximum Annual Rate: 7.5 pt/A. DO NOT exceed 1.9 lb ai/A/year of paraquat-containing products for preplant/preemergence uses. DO NOT exceed 1.9 lb ai/A/year of paraquat-containing products for all combined uses. DO NOT make more than 2 preemergence applications per year. DO NOT apply this product by air. DO NOT allow livestock to graze in treated areas. Pre-Harvest Interval (PHI): NA. 		

©2019 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company.

24(c) Registrant:
 Syngenta Crop Protection, LLC
 P. O. Box 18300
 Greensboro, NC 27419-8300

Label Code: MS1431044AA0519