

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS ONLY– NOT TO BE
USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A
CERTIFIED APPLICATOR.**

syngenta.

INTIRA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TENNESSEE

**Gramoxone® SL 2.0
For Weed Control in Row Middles of Lima Bean and Southern Pea**

**EPA Reg. No. 100-1431
EPA SLN No. TN-210003**

**This label expires and must not be distributed or used in accordance with this SLN
registration after June 2, 2026**

**KEEP OUT OF REACH OF CHILDREN /
MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.**



**DANGER / POISON
PELIGRO / VENENO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)
See additional precautionary statements and directions for use inside booklet.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

Target Weed	Rate pt/A	Application Timing	Use Directions
Palmer amaranth, other annual grass and broadleaf weeds	2.0-3.0	Postemergence (Directed with Hooded sprayer)	<p>Apply as a post-directed spray between the rows using a hooded sprayer. Apply when weeds are 2-4 inches tall and actively growing, larger weeds may not be adequately controlled. Use the higher rate on larger or hard to control weeds. Apply in a minimum of 10 gal/A of water.</p> <p>Use nozzles that provide uniform coverage within the treated area. Keep hoods on these sprayers adjusted to protect desirable vegetation. Operate equipment in a manner that avoids bouncing or raising the hoods off the ground. Tractor operating speed must be adjusted to avoid bouncing of the spray hoods.</p> <p>Add either: 1) a nonionic surfactant containing 80% or more surface-active agent at 0.25% v/v (2 pt/100 gal) of the finished spray volume OR 2) Crop Oil Concentrate or Methylated Seed Oil containing 15-20% approved emulsifier at 1.0% v/v (1 gal/100 gal) of the finished spray volume.</p> <p>A second application may be made, allow a minimum of 7 days between applications.</p>
Precaution: <ul style="list-style-type: none"> Spray should not contact green stems, foliage, or pods, as injury may result. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Maximum Single Application Rate: 3.0 pt/A REI: 24 hours Minimum Application Interval: 7 days Maximum Annual Rate: 6.0 pt/A DO NOT make more than 2 applications per year Pre-Harvest Interval (PHI): 21 days 			

©2021 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company,

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: TN1431032AA0521