

**RESTRICTED USE PESTICIDE  
DUE TO ACUTE TOXICITY  
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS ONLY– NOT TO BE  
USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A  
CERTIFIED APPLICATOR.**

**syngenta.**

**INTIRA**

## **Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TENNESSEE**

**Gramoxone® SL 2.0  
For Weed Control in Row Middles of Soybean, Immature Seed (Edamame)**

**EPA Reg. No. 100-1431  
EPA SLN No. TN-210004**

**This label expires and must not be distributed or used in accordance with this SLN  
registration after June 2, 2026**

**KEEP OUT OF REACH OF CHILDREN /  
MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.**



**DANGER / POISON  
PELIGRO / VENENO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)  
See additional precautionary statements and directions for use inside booklet.

### **DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

Target Weed	Rate pt/A	Application Timing	Use Directions
Palmer amaranth, other annual grass and broadleaf weeds	2.0-3.0	Postemergence (Directed with Hooded sprayer)	<p>Apply <b>prior to pod-set</b> as a post-directed spray between the rows using a hooded sprayer. Apply when weeds are 2-4 inches tall and actively growing, larger weeds may not be adequately controlled. Use the higher rate on larger or hard to control weeds. Apply in a minimum of 10 gal/A of water.</p> <p>Use nozzles that provide uniform coverage within the treated area. Keep hoods on these sprayers adjusted to protect desirable vegetation. Operate equipment in a manner that avoids bouncing or raising the hoods off the ground. Tractor operating speed must be adjusted to avoid bouncing of the spray hoods.</p> <p>Add either: 1) a nonionic surfactant containing 80% or more surface-active agent at 0.25% v/v (2 pt/100 gal) of the finished spray volume <b>OR</b> 2) Crop Oil Concentrate or Methylated Seed Oil containing 15-20% approved emulsifier at 1.0% v/v (1 gal/100 gal) of the finished spray volume</p> <p>A second application may be made, allow a minimum of 7 days between applications.</p>
<b>Precaution:</b> <ul style="list-style-type: none"> <li>Spray should not contact green stems, foliage, or pods, as injury may result.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ol style="list-style-type: none"> <li><b>Maximum Single Application Rate:</b> 3.0 pt/A</li> <li><b>REI:</b> 24 hours</li> <li><b>Minimum Application Interval:</b> 7 days</li> <li><b>Maximum Annual Rate:</b> 6.0 pt/A</li> <li><b>DO NOT</b> make more than 2 applications per year</li> <li><b>DO NOT</b> apply after pod-set</li> <li><b>Pre-Harvest Interval (PHI):</b> 45 days</li> </ol>			

©2021 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company,

24(c) Registrant:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, NC 27419-8300

Label Code: TN1431033AA0521