
SUPPLEMENTAL LABELING

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

Syngenta Crop Protection, LLC

P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1431A-S2 1211

Gramoxone® SL 2.0

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide
Grasses – (Kentucky Bluegrass grown for seed only) For Use Only in the State of Minnesota

Active Ingredient:

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride) 30.1%

Other Ingredients: 69.9%

Total: 100.0%

Contains 2.0 pounds paraquat cation per gallon as 2.762 pounds salt per gallon.
Contains alerting agent (odor), emetic, dye.

EPA Reg. No. 100-1431

KEEP OUT OF REACH OF CHILDREN.



**DANGER / POISON
PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- **NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS.**
- **IF SWALLOWED, TAKE IMMEDIATE ACTION AS PRESCRIBED IN FIRST AID STATEMENT. SYMPTOMS ARE PROLONGED AND PAINFUL.**
- **DO NOT USE OR STORE IN OR AROUND THE HOME.**
- **DO NOT REMOVE CONTENTS EXCEPT FOR IMMEDIATE USE.**
- **THE ODOR OF THIS PRODUCT IS FROM THE ALERTING AGENT WHICH HAS BEEN ADDED, NOT FROM PARAQUAT.**

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using GRAMOXONE SL 2.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

| Crop | Maximum Number of Applications | Use Pattern | Gramoxone SL 2.0 Rate/Acre | Minimum Total Spray per Acre | Grazing or Preharvest Interval | Precautions, Restrictions and Comments |
|--|--------------------------------|---|----------------------------|--------------------------------------|--------------------------------|--|
| GRASSES (Kentucky Bluegrass grown for seed only) For Use Only in the state of Minnesota | 2 | Post Harvest Desiccation to facilitate burning | 1.5-2.25 pts. | Ground: 10 gals. Air: 5 gals. | - | <ul style="list-style-type: none"> • For use on Kentucky bluegrass seed production fields in Minnesota as a post-harvest application to facilitate burning. • Add a nonionic surfactant at a rate of 1 pint per 100 gals. (75% or greater surface active ingredient) or 2 pints per 100 gallons (50-74% surface active ingredient). • Make application 1-4 weeks after harvest and wait 3-7 days before burning fields. • Do not graze or feed hay from treated fields prior to burning. • Do not apply this product through any type of irrigation system. |

Gramoxone® SL 2.0 and the Syngenta logo trademark of a Syngenta Group Company

©2012 Syngenta

GRAMOXONE SL 2.0 1431A-S2 1211 – tb – 02/08/12