

**SUPPLEMENTAL LABELING**

**RESTRICTED USE PESTICIDE  
DUE TO ACUTE TOXICITY**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

**Syngenta Crop Protection, LLC**

P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1431A-S6 1211**

**Gramoxone® SL 2.0**

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide  
Sugarcane – For Use Only in the State of Louisiana

Active Ingredient:

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride) ..... 30.1%

Other Ingredients: ..... 69.9%

Total: ..... 100.0%

Contains 2.0 pounds paraquat cation per gallon as 2.762 pounds salt per gallon.  
Contains alerting agent (odor), emetic, dye.

**EPA Reg. No. 100-1431**

**KEEP OUT OF REACH OF CHILDREN.**



**DANGER / POISON  
PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- **NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS.**
- **IF SWALLOWED, TAKE IMMEDIATE ACTION AS PRESCRIBED IN FIRST AID STATEMENT. SYMPTOMS ARE PROLONGED AND PAINFUL.**
- **DO NOT USE OR STORE IN OR AROUND THE HOME.**
- **DO NOT REMOVE CONTENTS EXCEPT FOR IMMEDIATE USE.**
- **THE ODOR OF THIS PRODUCT IS FROM THE ALERTING AGENT WHICH HAS BEEN ADDED, NOT FROM PARAQUAT.**

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using GRAMOXONE SL 2.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## DIRECTIONS FOR USE

Crop	Maximum Number of Applications	Use Pattern	Gramoxone SL 2.0 Rate/Acre	Minimum Total Spray per Acre	Grazing or Preharvest Interval	Precautions, Restrictions and Comments
<p><b>SUGARCANE</b></p> <p>For Use Only in the State of Louisiana</p>	<p style="text-align: center;">2</p>	<p style="text-align: center;">Broadcast</p> <p style="text-align: center;">Prior to sugarcane emergence</p> <p style="text-align: center;">Or</p> <p style="text-align: center;">When sugarcane has no more than 4 leaves</p>	<p style="text-align: center;">1.9-3.75 pts.</p>	<p style="text-align: center;">Ground: 10 gals.</p> <p style="text-align: center;">Air: 5 gals.</p>	<p style="text-align: center;">30 days</p>	<ul style="list-style-type: none"> <li>• For control of annual ryegrass and other weeds prior to or after crop emergence (but only when sugarcane has no more than four leaves) in sugarcane in Louisiana.</li> <li>• For improved burndown and residual control, tank mix Gramoxone SL 2.0 with atrazine or diuron (Karmex or Direx).</li> <li>• For best results, apply when weeds and grasses are succulent and weed growth is 1 to 6 inches.</li> <li>• Use higher rates for larger weeds.</li> <li>• Do not exceed 6 pts. Gramoxone SL 2.0 per acre per season.</li> <li>• Failure to add a nonionic surfactant or crop oil concentrate will result in reduced performance of Gramoxone SL 2.0. Add either: 1) a nonionic surfactant containing 75% or more surface-active agent at 0.125% v/v (1 pt. per 100 gals), <b>OR</b> 2) add a nonionic surfactant containing 50-74% surface-active agent at 0.25% v/v (2 pts. per 100 gals.), of the finished spray volume for ground applications. For aerial applications, add 1) a nonionic surfactant at 0.25% v/v (2pts. per 100 gals.) of the finished spray volume, <b>OR</b> 2) A crop oil concentrate (nonphytotoxic) containing 15-20% approved emulsifier, at 1.0% v/v (1gal. per 100 gals.) of the finished spray volume for ground applications. For aerial applications, add 1 pt. of crop oil concentrate per acre.</li> <li>• Do not apply this product through any type of irrigation system.</li> <li>• Do not graze treated areas of feed treated forage to livestock.</li> </ul>

Gramoxone® SL 2.0 and the Syngenta logo trademark of a Syngenta Group Company

©2012 Syngenta

GRAMOXONE SL 2.0 1431A-S6 1211 – tb – 02/08/12