#### SUPPLEMENTAL LABELING

# RESTRICTED USE PESTICIDE DUE TO ACUTE TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

## Syngenta Crop Protection, LLC

P. O. Box 18300 Greensboro, North Carolina 27419-8300 SCP 1431A-S9 1211

#### Gramoxone® SL 2.0

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide Trees and Vines (Grapes) – For Use Only in the State of California

#### **Active Ingredient:**

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride)	30.1%
Other Ingredients:	69.9%
Total:	100.0%

Contains 2.0 pounds paraquat cation per gallon as 2.762 pounds salt per gallon. Contains alerting agent (odor), emetic, dye.

EPA Reg. No. 100-1431

### KEEP OUT OF REACH OF CHILDREN.



# DANGER / POISON PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS.
- IF SWALLOWED, TAKE IMMEDIATE ACTION AS PRESCRIBED IN FIRST AID STATEMENT. SYMPTOMS ARE PROLONGED AND PAINFUL.
- DO NOT USE OR STORE IN OR AROUND THE HOME.
- DO NOT REMOVE CONTENTS EXCEPT FOR IMMEDIATE USE.
- THE ODOR OF THIS PRODUCT IS FROM THE ALERTING AGENT WHICH HAS BEEN ADDED, NOT FROM PARAQUAT.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using GRAMOXONE SL 2.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



# **DIRECTIONS FOR USE**

	Maximum		_		Grazing	
	Number		Gramoxone	Minimum	or	
	of		SL 2.0	Total Spray	Preharvest	
Crop	Applications	Use Pattern	Rate/Acre	per Acre	Interval	Precautions, Restrictions and Comments
TREES AND VINES GRAPES	1	Broadcast aerial application	1.1-1.9 pts.	Air: 10 gals.		Gramoxone SL 2.0 can be applied by air for control of emerged annual weeds in grapes (dormant application).
For Use Only in		to dormant vines				Apply when weeds are succulent and growth is from 1 to 6 inches high.
the State of California						Add a nonionic surfactant containing 50% surface-active agent at 1-2 pts. per 100 gallons of the finished spray volume.
						Do not apply more than once per season.
						Do not enter treated field within 24 hours after application.
						Do not apply if internal tissues of the buds are exposed as injury to the developing buds will result.

Gramoxone® SL 2.0 and the Syngenta logo trademark of a Syngenta Group Company

©2012 Syngenta

GRAMOXONE SL 2.0 1431A-S9 1211 - tb - 02/08/12