RESTRICTED USE PESTICIDE DUE TO ACUTE TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS— NOT TO BE USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR.



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MONTANA

Gramoxone® SL 3.0

For Use as a Harvest Aid in Alfalfa Grown for Seed

EPA Reg. No. 100-1652 EPA SLN No. MT-21-0001

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2026

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) See additional precautionary statements and directions for use inside booklet

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

Alfalfa Grown for Seed

| Crop | Use Pattern | Rate Per Treated Acre | Minimum Spray Volume gal/acre | Preharvest Interval (Days) |
|--|-------------|--------------------------|----------------------------------|-------------------------------|
| Alfalfa | Broadcast | 1.7 – 2.7 pt | Ground: 20-25 gal | See precautions below |
| Desiccation of alfalfa to facilitate harvest of alfalfa seed | | | Air: 5-10 gal | |

Always Add One of the Following

Add a nonionic surfactant containing 80% or more surface-active agent at 0.25% v/v (2 pint per 100 gallons) of the finished spray volume for ground applications. For aerial applications, add a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons) of the finished spray volume.

Precautions, Restrictions, and Comments

- Restricted Entry Interval (REI): 48 hours
- DO NOT apply Gramoxone SL 3.0 if grain crops growing in directly adjacent fields have not been harvested.
- DO NOT apply more than 2 applications per year.
- DO NOT exceed 1.0 lb ai/A/year of paraquat containing products for all combined uses
- DO NOT harvest until at least 4 days after application.
- DO NOT apply when weather conditions favor drift from treated areas.
- DO NOT apply by ground equipment within 25 ft or by air within 75 ft of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries; and commercial fish ponds.
- For use only on fields in production of alfalfa seed. Not for use on fields producing alfalfa for livestock feed. No portion of the treated field, including seed, seed screenings, hay forage, or stubble, may be used for human or animal feed. Screenings from alfalfa seed processing are prohibited from feed channels. All Gramoxone SL 3.0 treated alfalfa seed screenings must be removed from the feed market.
- DO NOT cut current year's treated alfalfa seed crop for hay or forage. DO NOT graze current year's treated alfalfa seed crops.
- Treated alfalfa seed is not to be used for sprouting. All alfalfa seed treated with Gramoxone SL 3.0 is to be tagged at processing plants, "NOT FOR HUMAN CONSUMPTION OR ANIMAL FEED". It shall be the grower's responsibility to notify the processing plants of any seed crop treated with Gramoxone SL 3.0.
- DO NOT apply this product through any type of irrigation system.

©2021 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant: Syngenta Crop Protection, LLC P. O. Box 18300 Greensboro, North Carolina 27419-8300

Label Code: MT1652051AA0821