

## SUPPLEMENTAL LABELING

### RESTRICTED USE PESTICIDE DUE TO ACUTE TOXICITY

TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR, EXCEPT THAT UNCERTIFIED PERSONS MAY TRANSPORT CONTAINERS AS PROVIDED UNDER DIRECTIONS FOR USE.

#### Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina, 27419-8300

SCP 1652A-S21A 0822

PARAQUAT DICHLORIDE

GROUP

22

HERBICIDE

#### Gramoxone® SL 3.0

#### HERBICIDE

EPA Reg. No. 100-1652

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

#### Active Ingredients:

Paraquat dichloride\* ..... 43.9%

Other Ingredients: ..... 56.1%

**Total:** ..... **100.0%**

\*CAS No. 4685-14-7

Gramoxone SL 3.0 is formulated as a liquid containing 3.0 pounds paraquat cation per gallon as 4.134 pounds paraquat dichloride per gallon.

Gramoxone SL 3.0 contains alerting agent (odor), emetic, and dye.

**KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.**



**DANGER / POISON**  
**PELIGRO / VENENO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Gramoxone SL 3.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**syngenta**

**DIRECTIONS FOR USE  
RESTRICTED USE PESTICIDE**

Potato, ME Only

Crops (Including all cultivars, varieties, and/or hybrids of these)			
Potato			
Target Weed	Rate pt/A	Application Timing	Use Directions
Refer to <b>Section 7.0.</b>	1.7	Postemergence Desiccation  5 days minimum application interval	<p>For use in <b>Maine Only</b> for desiccation of potato plants to control late blight disease.</p> <p><b>This is a CROP DESTROY use. Crop must be destroyed and tubers not allowed to enter channels of trade. All tubers must be disked-in after desiccation</b></p> <p>Apply as a ground broadcast over-the-top application of infected plants in a minimum of 20 gal/A of spray volume.</p> <p>Add a nonionic surfactant at 1.0 quart per 100 gallons of spray mix.</p> <p>Make a second application if necessary to obtain additional desiccation where vine growth is dense.</p> <p>See <b>Section 4.4.4</b> of the Gramoxone SL 3.0 label for adjuvant recommendations.</p>
<b>Tank Mix Options:</b> <ul style="list-style-type: none"> <li>Gramoxone SL 3.0 can be tank mixed with other herbicides labeled for use on potatoes applied by the same methods and at the same timings. Refer to <b>Section 4.4.2</b> for tank-mix precautions in the Gramoxone SL 3.0 label.</li> </ul>			
<b>Resistance Management:</b> <ul style="list-style-type: none"> <li>Refer to <b>Section 3.1.</b></li> </ul>			
USE RESTRICTIONS			
<ol style="list-style-type: none"> <li>Refer to <b>Section 6.1</b> for additional product use restrictions.</li> <li><b>Maximum Single Application Rate:</b> 1.7 pt/A</li> <li><b>REI:</b> 48 hours</li> <li><b>Minimum Application Interval:</b> 5 days</li> <li><b>Maximum Annual Rate:</b> 3.4 pt/A               <ol style="list-style-type: none"> <li><b>DO NOT</b> exceed 1.2 lb cation/A/year of paraquat-containing products.</li> </ol> </li> <li><b>DO NOT</b> make more than 2 application per year.</li> <li><b>This is a CROP DESTROY use. Crop must be destroyed and tubers not allowed to enter channels of trade. All tubers must be disked-in after desiccation.</b></li> <li><b>DO NOT</b> apply this product through any type of irrigation system.</li> <li><b>DO NOT</b> graze livestock in treated area.</li> <li><b>Pre-Harvest Interval (PHI):</b> Harvest crop at normal maturity.</li> </ol>			

Gramoxone® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2022 Syngenta

For non-emergency (e.g., current product information), call  
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina, 27419-8300