SUPPLEMENTAL LABELING

RESTRICTED USE PESTICIDE DUE TO ACUTE TOXICITY

TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR, EXCEPT THAT UNCERTIFIED PERSONS MAY TRANSPORT CONTAINERS AS PROVIDED UNDER DIRECTIONS FOR USE.

Syngenta Crop Protection, LLC

P. O. Box 18300 Greensboro, North Carolina, 27419-8300 SCP 1652A-S11A 0822

PARAQUAT DICHLORIDE GROUP 22 HERBICIDE

Gramoxone® SL 3.0 HERBICIDE

EPA Reg. No. 100-1652

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

Active Ingredients:

Paraquat dichloride* 43.9%
Other Ingredients: 56.1%

Total:

100.0%

*CAS No. 4685-14-7

Gramoxone SL 3.0 is formulated as a liquid containing 3.0 pounds paraquat cation per gallon as 4.134 pounds paraquat dichloride per gallon.

Gramoxone SL 3.0 contains alerting agent (odor), emetic, and dye.

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)
See additional precautionary statements and directions for use inside booklet.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Gramoxone SL 3.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE RESTRICTED USE PESTICIDE

Grapes, CA Only

Crops (Including all cultivars, varieties, and/or hybrids of these)

Grapes

Target Weed	Rate pt/A	Application Timing	Use Directions
Refer to Section 7.0.	0.7 – 1.3	Dormant Application	For use only for use in California Only as a broadcast aerial application to dormant vines for control of emerged annual weeds.
			Apply in a minimum of 10 gal/A of spray volume when weeds are succulent and growth is 1 to 6".
			Gramoxone SL 3.0 can be applied by air for control of emerged annual weeds in grapes (dormant application).
			Add a nonionic surfactant containing 50% surface-active agent at 1-2 pt per 100 gallons of the finished spray volume.
			See Section 4.4.4 of the Gramoxone SL 3.0 label for adjuvant recommendations.

Tank Mix Options:

 Gramoxone SL 3.0 can be tank mixed with other herbicides labeled for use on grapes applied by the same methods and at the same timings. Refer to Section 4.4.2 for tank-mix precautions in the Gramoxone SL 3.0 label.

Resistance Management:

• Refer to **Section 3.1.**

Precaution:

DO NOT apply if internal tissues of the buds are exposed as injury to the developing buds will
result.

USE RESTRICTIONS

- 1. Refer to **Section 6.1** for additional product use restrictions.
- 2. Maximum Single Application Rate: 1.3 pt/A
- 3. **REI:** 48 hours
- 4. Minimum Application Interval: Not Applicable
- 5. Maximum Annual Rate: 1.3 pt/A
 - a. **DO NOT** exceed 0.5 lb cation/A/year of paraquat-containing products.
- 6. **DO NOT** make more than one application per year.
- 7. Pre-Harvest Interval (PHI): Harvest crop at normal harvest maturity.

Gramoxone®, and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2022 Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.

Manufactured for: Syngenta Crop Protection, LLC P.O. Box 18300 Greensboro, North Carolina, 27419-8300