

SUPPLEMENTAL LABELING

RESTRICTED USE PESTICIDE

DUE TO ACUTE TOXICITY

TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR.

PARAQUAT DICHLORIDE

GROUP

22

HERBICIDE

HELMQUAT 3SL

A Weed, Grass and Harvest Aid Desiccant/ Defoliant Herbicide

PEANUTS

Recirculating Rope Wick or Carpet Roller Equipment

Suppression or control of Glyphosate-Resistant Palmer Amaranth [and Florida Beggarweed]

Active Ingredient	% BY WT.
Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride).....	43.8%
Other Ingredients.....	56.2%
Total.....	100.0%

Contains 3.0 pounds paraquat cation per gallon as 4.14 pounds paraquat dichloride salt per gallon.

Contains stench (odor) and emetic.

EPA Reg. No. 74530-48

Expires: December 31, 2022

KEEP OUT OF REACH OF CHILDREN/MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

DANGER/PELIGRO

POISON/VENENO



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

CORROSIVE TO SKIN AND EYES	CORROSIVO PARA LA PIEL Y LOS OJOS.
NEVER TRANSFER THIS PRODUCT INTO FOOD OR BEVERAGE CONTAINERS OR CONTAINERS NOT EXPLICITLY INTENDED FOR PESTICIDES	NUNCA TRANSFERIRA ESTE PRODUCTO A RECIPIENTES PARA COMIDA O DE BEBIDAS O RECIPIENTES NO EXPLICITAMENTE PREVISTOS PARA PLAGUICIDAS.
READ ENTIRE LABEL PRIOR TO USING THIS PRODUCT	LEA LA ETIQUETA COMPLETA ANTES DE USAR ESTE PRODUCTO.
IN THE CASE OF AN ACCIDENT, SEEK IMMEDIATE MEDICAL ATTENTION. SYMPTOMS ARE PROLONGED, PAINFUL, AND CAN BE FATAL	EN CASO DE ACCIDENTE, BUSQUE ATENCIÓN MÉDICA INMEDIATA. LOS SÍNTOMAS SON PROLONGADOS, DOLOROSOS, Y PUEDEN SER MORTALES.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING.

This label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE PRODUCT LABEL BOOKLET MUST BE FOLLOWED, INCLUDING STATEMENTS PERTAINING TO THE WORKER PROTECTION STANDARDS, ON THE EPA REGISTERED LABEL. BEFORE USING HELMQUAT 3 SL AS PERMITTED ACCORDING TO THIS SUPPLEMENTAL LABEL, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL AND THE ATTACHED PESTICIDE PRODUCT CONTAINER.

Read the label affixed to the container for HELMQUAT 3SL before applying.

Use of HELMQUAT 3SL according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for HELMQUAT 3SL.

CROP	RATE PER ACRE	COMMENTS
PEANUTS AR, FL, GA, MS, OK, NC, SC [other states may be added as deemed appropriate] Suppression of glyphosate- resistant Palmer Amaranth and Florida Beggarweed	Mix 1 part HELMQUAT 3SL with 1-1.5 parts water to prepare a 40-50% solution. Add non-ionic surfactant containing 75% or more surface-active agent at a minimum of 0.25% v/v (2 pints/100 gal) of finished volume. Apply up to 2 pints/acre of the herbicide-water mixture. RESTRICTIONS <ul style="list-style-type: none"> • Do not apply more than 1 pints/acre (0.25 lbs. a.i./acre) of HELMQUAT 3SL using the recirculating wick or carpet roller application method. • Do not exceed 0.5 lb. a.i./acre total for all HELMQUAT 3SL applications to peanut. • Make application of HELMQUAT 3SL as early as possible, and not less than 30 days before harvest to control weeds that may interfere with harvesting operations. • Do not allow livestock to graze in treated areas. • Do not feed hay or threshings from treated fields to livestock. 	Apply through a recirculating rope wick or carpet roller equipment for suppression or control of glyphosate-resistant Palmer Amaranth [and Florida Beggarweed] to prevent seed production. In order to prevent seed production, an application must be made prior to formation of viable seed. Make application at least 6 inches above the peanut canopy. Set application equipment to avoid dripping. Low ground speed enhances coverage which leads to improved weed control. Use ground speeds of 5 mph or less. Delaying application to late afternoon or early evening will enhance control of large weeds. Follow application equipment manufacturer's directions.

This HELM Supplemental Label is to expire on December 31, 2022 and should not be used after that date.

NOTICE

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Manufactured for:



HELM AGRO US, Inc.
 401 E. Jackson St., Suite 1400
 Tampa, FL 33602
info@helmagro.com
[Phone: 813-621-8846](tel:813-621-8846)