

# SUPPLEMENTAL LABELING

## RESTRICTED USE PESTICIDE

### DUE TO ACUTE TOXICITY

TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR.

PARAQUAT DICHLORIDE

GROUP

22

HERBICIDE

## HELMQUAT 3SL

A Weed, Grass and Harvest Aid Desiccant/ Defoliant Herbicide

### VEGETABLES - FLORIDA

#### Cabbage & Lettuce

Active Ingredient	% BY WT.
Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride).....	43.8%
Other Ingredients.....	56.2%
Total.....	100.0%

Contains 3.0 pounds paraquat cation per gallon as 4.14 pounds paraquat dichloride salt per gallon.

Contains stench (odor) and emetic.

EPA Reg. No. 74530-48

Expires: December 31, 2022

KEEP OUT OF REACH OF CHILDREN/MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

**DANGER/PELIGRO**

**POISON/VENENO**



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

CORROSIVE TO SKIN AND EYES	CORROSIVO PARA LA PIEL Y LOS OJOS.
NEVER TRANSFER THIS PRODUCT INTO FOOD OR BEVERAGE CONTAINERS OR CONTAINERS NOT EXPLICITLY INTENDED FOR PESTICIDES	NUNCA TRANSFEIERA ESTE PRODUCTO A RECIPIENTES PARA COMIDA O DE BEBIDAS O RECIPIENTES NO EXPLÍCITAMENTE PREVISTOS PARA PLAGUICIDAS.
READ ENTIRE LABEL PRIOR TO USING THIS PRODUCT	LEA LA ETIQUETA COMPLETA ANTES DE USAR ESTE PRODUCTO.
IN THE CASE OF AN ACCIDENT, SEEK IMMEDIATE MEDICAL ATTENTION. SYMPTOMS ARE PROLONGED, PAINFUL, AND CAN BE FATAL	EN CASO DE ACCIDENTE, BUSQUE ATENCIÓN MÉDICA INMEDIATA. LOS SÍNTOMAS SON PROLONGADOS, DOLOROSOS, Y PUEDEN SER MORTALES.

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING.

This label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE PRODUCT LABEL BOOKLET MUST BE FOLLOWED, INCLUDING STATEMENTS PERTAINING TO THE WORKER PROTECTION STANDARDS, ON THE EPA REGISTERED LABEL. BEFORE USING HELMQUAT 3 SL AS PERMITTED ACCORDING TO THIS SUPPLEMENTAL LABEL, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL AND THE ATTACHED PESTICIDE PRODUCT CONTAINER.

Read the label affixed to the container for HELMQUAT 3SL before applying.

Use of HELMQUAT 3SL according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for HELMQUAT 3SL.

CROP	RATE PER ACRE	COMMENTS
<b>VEGETABLES:</b> <b>Cabbage</b> <b>Lettuce</b> [FL only]	0.8-1.27 pts. <b>For Cabbage</b> add either Non-ionic surfactant at 0.25% v/v (or 32 fl. oz.) of finished volume, Or Crop Oil Concentrate (non-phytotoxic) containing 15-20% approved emulsifier, at 1.0 v/v (1 gallon per 100 gallons) of the finished spray volume.  <b>For Lettuce</b> add Non-ionic surfactant at 0.25% v/v (or 32 fl. oz.) of finished volume.	HELMQUAT 3SL may be used as a postemergent directed spray – shielded application in a minimum of 40 gallons of spray for control of emerged weeds in rows middles in cabbage (including tight head Chinese) and lettuce. Apply with conventional ground equipment directing spray between the rows and using shields to prevent spray contact with crop plants. Use properly configured precision directed spray equipment adjusted to prevent spray contact with crop plants. Apply when weeds are succulent and weed growth is less than 6" in height. Weeds emerging after application will not be controlled.
<b>RESTRICTIONS</b> <ul style="list-style-type: none"> <li>• Do not make more than 3 applications per season.</li> <li>• Do not allow spray to contact crop as damage or excessive residues may result.</li> <li>• Do not apply via chemigation.</li> <li>• Do not allow animals to graze in treated areas.</li> <li>• Do not apply where HELMQUAT 3SL (or other paraquat containing product) has been used as a preplant preemergence spray.</li> <li>• Strip outer leaves at the time of harvest.</li> </ul>		

**This HELM Supplemental Label is to expire on December 31, 2022 and should not be used after that date.**

### NOTICE

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Manufactured for:



HELM AGRO US, Inc.  
 401 E. Jackson St., Suite 1400  
 Tampa, FL 33602  
[info@helmagro.com](mailto:info@helmagro.com)  
[Phone: 813-621-8846](tel:813-621-8846)