

HingeTM EPA Reg. No. 83100-40-83979 2(ee) Recommendation

HING PLUS CHATEAU HERBICIDE SW OR OXYSTAR 4L* OR OXYSTAR 2E** FOR IMPROVED WEED CONTROL IN THE STATES OF ARIZONA, CALIFORNIA, IDAHO, OREGON, UTAH, AND WASHINGTON IN LABELED TREE AND VINE CROPS.

This recommendation is permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions for use, precautions and limitations on the product label attached to the container for Hinge herbicide. Users must have this recommendation in their possession at the time of use.

Refer to the individual product labels for products listed for precautionary statements, restrictions, use rates, approved uses, and a list of weeds controlled.

Hinge applied at the rate of 4.0 oz product per acre can be tank mixed with Chateau Herbicide SW at 6.0 to 12.0 ounces per acre, or Oxystar 4L* at 1.5 to 3.0 pints per acre, or Oxystar 2E** at 3.0 to 6.0 pints per acre to help improve overall weed control in labeled crops.

In order to obtain the best residual weed control from Hinge, it is important to use a postemergence/contact herbicide such as glyphosate or other registered postemergence herbicides to control any emerged weeds.

Do not apply more than 4.0 oz Hinge per year on a broadcast acre basis. When applied as a banded treatment (50% band or less), Hinge may be applied twice per year. A minimum of 30 days must be allowed between applications of Hinge.

Weed control is limited to weeds on the Hinge label.

Effective Date: 06.27.16

Hinge, Oxystar 4L, and Oxystar 2E are registered trademark of Albaugh, LLC.

Chateau Herbicide SW is a registered trademark of Valent U.S.A. Corporation.

- * Not registered in Washington
- ** Not registered in Utah

