



HoloPhos™

Guaranteed analysis: 0-7-0

Available Phosphate (P₂O₅).....7%
Calcium (Ca)7%

Derived from: Rock phosphate

ALSO CONTAINS NONPLANT FOOD INGREDIENTS:

0.1% humic acid (derived from leonardite)

- 1 gallon (3.78L) 55 gallon (208.2L)
- 5 gallon (18.92L) 275 gallon (1041L)
- 30 gallon (113.56L)

Density: 10.4 lbs. per gallon at 68° F

Volume: 1 gallon (3.78L)

Information regarding the contents
and levels of metals in this product is
available on the internet at

<http://www.aapfco.org/metals.htm>

This product is intended for use according to an approved organic system plan. All materials comply with the USDA National Organic Program (NOP) rules and other international bio-organic regulations for certified organic crop production. Verify with your certifier before using.

Application rates and timing

This material is intended for foliar application and should be used in accordance with manufacturer's instructions.

Product Purpose: This product provides phosphorus for overall plant health and nutrition.

Fruits and Vegetables: As a foliar and/or in irrigation, apply at a rate of 1-2 quarts per acre, every 7-14 days as part of a comprehensive fertility program.

Broad Acre Crops: For use as a row starter, or as a side dress, can be applied at 4-8 quarts per acre, depending on soil analysis. As a foliar, apply 2-4 quarts per acre at appropriate plant growth stages.

Maximum target particle size contained in this product is 100 µm. Use appropriate spray screens.

Handling: Store in cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95° F. Shake or stir before use, especially when near empty to dissolve the residue on the bottom of the drum/bottle. Do not pre-mix or store in diluted form. Always use product within 8 hours of diluting. All foliar sprays perform best when applied early morning or late afternoon.

When tank mixing: A jar test is recommended: Prepare a small amount of the proposed mix and allow it to stand in a glass jar over night observing for signs of incompatibility. Ideally, a sample of the test mix should then be sprayed on a small area of the target crop and observed for 3 to 4 days for any phytotoxic effects. Agitation is the key to usability of this product. AEA products involve tiny particles which are far too small to clog filters, however if they are inadequately diluted or agitated before and during application there is a possibility of problems with spray equipment. Consider using an eductor nozzle to agitate prior to use. Compatibility and/or performance cannot be guaranteed when combined with other products.

Disclaimer: AEA products are sold "as is" and all warranties, express or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.



Advancing Eco Agriculture, LLC
4551 Parks West Road, Middlefield, OH 44062
www.advancingecoag.com • 800-495-6603