

FIFRA 2(ee) Recommendation



EPA Reg. No. 91197-3-92488

ADDING CONTROL of DOWNY MILDEWS, PESTALOTIOPSIS, and POWDERY MILDEWS on ALL LABELED CROPS,

and

ADDING CONTROL of BLACK ROT, BOTRYOSPHAERIA, CHARCOAL ROT, GREASY SPOT, GUMMY STEM BLIGHT, MELANOSE, MONILINIA, PHOMOPSIS, SHOT HOLE, and SPUR BLIGHT on SELECTED CROPS

ADDING SUPPRESSION of FIRE BLIGHT, RUST, and SCAB on SELECTED CROPS

CLARIFYING ACCEPTABILITY of IN-FURROW, DRENCH OR TRANSPLANT STARTER SOLUTION USE of HOWLER on ALL LABELED CROPS

FOR USE AND DISTRIBUTION ONLY IN THE STATE OF CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. All applicable directions, restrictions, precautions, and Conditions of Sale or Warranty on the EPA registered label are to be followed. Refer to the container label for additional instructions. Always read and follow label directions.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

ADDING CONTROL of DOWNY MILDEWS, PESTALOTIOPSIS, and POWDERY MILDEWS on ALL LABELED CROPS, and ADDING CONTROL of VARIOUS DISEASES on SELECTED CROPS:

Disease Control	Crop
Alternaria	All labeled crops
(Alternaria spp.)	All labeled crops
Black rot	Grapes
(Guignardia spp.)	•
	Berries and Small Fruits
Botryosphaeria	O(and Emilia (analysis and amount)
(Botryosphaeria spp.)	Stone Fruits (excluding plums and prunes)
	Tree Nuts
Downy mildews	
(Pseudoperonospora spp.,	
Plasmopara spp., Sclerophthora spp.,	All labeled crops
Phytophthora spp., Peronospora spp.,	
Plasmopara spp., and Bremia spp.)	
Greasy spot	Citrus
(Mycosphaerella spp.)	
Gummy stem blight	Cucurbit Vegetables
(Stagonosporopsis spp.)	
Melanose	Citrus
(Diaporthe spp.)	Daming and Oncell Enviro
	Berries and Small Fruits
Monilinia	Stone Fruits (excluding plums and prunes)
(<i>Monilinia</i> spp.)	Storie Fruits (excluding plums and prunes)
	Tree Nuts
Pestalotiopsis	All laboled crops
(Pestalotiopsis spp.)	All labeled crops
Phomopsis	Berries and Small Fruits
(<i>Phomopsis</i> spp.)	
(Grapes



Disease Control	Crop
Powdery mildews	
(Podosphaera spp., Erysiphe spp.,	
Sphaerotheca spp., Leveillula spp.,	All labeled crops
Microsphaera spp., Phyllactinia spp.,	
Uncinula spp., Oidium spp.)	
Shot hole	Stone Fruits (excluding plums and prunes)
(Wilsonomyces spp.)	
(Wilsonomyces spp.)	Tree Nuts
Spur blight	Raspberry
(<i>Didymella</i> spp.)	ιλαορυστιγ

For control of charcoal rot (*Macrophomina* spp.) on strawberries:

Apply 2.5-7.5 lbs/A as a foliar spray or via chemigation.

ADDING SUPPRESSION OF FIRE BLIGHT, RUST, and SCAB on SELECTED CROPS:

Disease Suppression	Crop
Fire blight	
(<i>Erwinia</i> spp.)	Apple
Scab	Apple
(Venturia spp.)	

For suppression of rust (*Puccinia* spp.) on small fruits:

Apply 2.5-7.5 lbs/A as a foliar spray or via chemigation. Apply in rotations or mixtures with other effective fungicides.

CLARIFYING ACCEPTABILITY OF IN-FURROW, DRENCH OR TRANSPLANT STARTER SOLUTION USE OF HOWLER on ALL LABELED CROPS:

For all labeled crops, Howler® fungicide may be applied as an in-furrow spray, drench or transplant starter solution at an application rate of 2.5-5 lbs Howler® fungicide/acre.

©2022 AgBiome Innovations, Inc. All Rights Reserved