ACCEPTED

Jan 29, 2021

Doc id: 571922

AGBIOME INNOVATIONS®

New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration

FIFRA 2(ee) Recommendation



ADDING SUPPRESSION OF ALTERNARIA, DOWNY MILDEWS, PESTALOTIOPSIS, AND POWDERY MILDEWS IN ALL LABELED CROPS CLARIFYING ACCEPTABILITY OF IN-FURROW, DRENCH OR TRANSPLANT STARTER SOLUTION USE OF HOWLER IN ALL LABELED CROPS EPA REG. No. 91197-3-92488

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. All applicable directions, restrictions, precautions, and Conditions of Sale or Warranty on the EPA registered label are to be followed. Refer to container label for additional instructions. Always read and follow label directions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

DIRECTIONS FOR USE

For Suppression of Alternaria, Downy Mildews, Pestalotiopsis, and Powdery Mildews

For suppression of Alternaria (Alternaria spp), Downy Mildews (Pseudoperonospora spp, Plasmopara spp, Sclerophthora spp, Phytophthora spp, Peronospora spp, Plasmopara spp, and Bremia spp), Pestalotiopsis (Pestalotiopsis spp), Powdery Mildews (Podosphaera spp, Erysiphe spp, Sphaerotheca spp, Leveillula_spp, Microsphaera spp, Phyllactinia spp, Uncinula spp, Oidium spp), in all labeled crops, apply Howler® Fungicide at 5 lb/acre. Howler® Fungicide can be applied by incorporation, drench, drip applications, in-furrow, foliar or chemigation.

For All Labeled Crops

For all labeled crops, Howler® Fungicide may be applied as an in-furrow spray, drench or transplant starter solution at an application rate of 5 - 15 lbs Howler® Fungicide/acre or 67 - 100 oz Howler® Fungicide/100 gal water.

© 2020 AgBiome Innovations®. All rights reserved.