

<b>FIRST AID:</b> Have product container or label with you when calling a poison control center or doctor or going for treatment. In case of poisoning, contact your Poison Control Center at 1-800-222-1222.
<b>If Swallowed:</b> Call poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to by poison control center or doctor. Do not give any liquid to a person. Do not give anything to an unconscious person.
<b>If in Eyes:</b> Hold eye open and rinse slowly and gently with plenty of water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call poison control center or doctor for treatment advice.
<b>If on Skin:</b> Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.
<b>If Inhaled:</b> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration by mouth-to-mouth, if possible. Call poison control center or doctor for treatment advice.
<b>Note To Physician:</b> This product contains petroleum distillates and may pose an aspiration pneumonia hazard.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION:** Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not store, use or spill near heat or open flame. Do not weld, heat or drill container. Residue may ignite with explosive violence if heated sufficiently.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical-resistance category selection chart. Mixers, loaders, applicators, flaggers and other handlers must wear: 1. Long-sleeved shirt and long pants. 2. Chemical-resistant gloves, made of barrier laminate or nitrile rubber or neoprene rubber or viton. 3. Shoes plus socks. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

**ENGINEERING CONTROLS**

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

**USER SAFETY RECOMMENDATIONS**

- Users should:
1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
  2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
  3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with this labeling.

Do not apply in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**CHEMIGATION PROHIBITION**

DO NOT apply this product through any type of irrigation system.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements of training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is: 1. Coveralls, 2. Chemical-resistant gloves made of any waterproof material, 3. Shoes plus socks.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter until sprays have dried.



# DORMANT OIL

ACTIVE INGREDIENT: Mineral Oil*	98.5%
INERT INGREDIENTS	1.5%
TOTAL	100.0%
*Contains Petroleum Distillates	
Minimum Unsulfonated Residue	93.5%

## KEEP OUT OF REACH OF CHILDREN

# CAUTION

SEE OTHER CAUTIONS ON SIDE PANEL

EPA REG. NO. 71058-2	NET CONTENTS:
EPA EST. NO. 47255 WA-001	_____ Gallons

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. **Pesticide Storage:** Keep container closed when not in use. Protect from freezing. Keep in shade. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal: Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container (greater than five gallons):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Refillable Container:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE), CALL INFOTRAC AT 1-800-535-5053.**

**SPRAY DRIFT MANAGEMENT**

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product. Wind Speed: Do not apply at wind speeds greater than 15 mph at the application site. Temperature Inversions: If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions. Droplet Size: Apply as a medium or coarser spray (ASABE standard 572), and the minimum volume mean diameter (VMD) for spinning atomizer nozzles.

**GROUND APPLICATION:** Apply using a nozzle height of no more than 4 feet above the ground or crop canopy. Applications must be consistent with the limitations for wind speed, temperature inversions, and droplet size indicated above.

**AIRBLAST APPLICATION:** For airblast application, turn off outward-pointing nozzles at row ends and when spraying outer row. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy. Applications must be consistent with the limitations for wind speed, temperature inversions, and droplet size indicated above.

**AERIAL APPLICATION:** Applications must be consistent with the limitations for wind speed, temperature inversions, and droplet size indicated above.

Release Height: For aerial application of agricultural products, do not release spray at a height more than 10 feet above the ground, the tops of crops, or above the orchard canopy.

Boom Length: The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Swath Adjustment: When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Leave at least one swath unsprayed at the downwind edge of the treated field.

**IAP DORMANT OIL** is narrow range petroleum spray oil designed for use on stone fruits, nut crops, pome fruits, and ornamentals as a foliar or dormant/delayed dormant application. Total coverage of all parts of plants is required for optimum results. Application rates for dilute sprays (greater than 150 gallons spray solution per acre) are listed as gallons of spray oil per 100 gallons of water per acre. Concentrate applications (usually from 20 to 150 gallons per acre), are based upon dilute spray conversion of rate per acre x highest volume applied per acre. Aerial applications should be considered only when weather conditions do not allow ground application. Apply specified amount of product by air in a minimum of 20 gallons per acre. Optimum spray gallonage per acre is based upon tree or plant size, planting density, and stage of growth.

**MIXING INSTRUCTIONS:** Fill spray tank with ½ the required amount of water. Add micronutrients and other compatible pesticides. If dry products are used, start agitation and mix thoroughly with water before adding oil to the spray tank. Add spray oil and continue with vigorous agitation. IAP Dormant Oil is formulated as a “fast break” spray oil, and will not stay in solution unless agitated constantly. Maintain agitation of the spray mixture until it has been applied. **DO NOT USE THIS MATERIAL IF IT DOES NOT EMULSIFY.**

**USE RESTRICTIONS**

- DO NOT COMBINE dormant spray oil with SULFUR or LIME SULFUR unless the mixture has been tested on the crop and the mixture has been shown to be safe. Oil and Sulfur or lime sulfur combinations SHOULD ONLY be used on DORMANT CROPS.
- DO NOT USE IAP DORMANT OIL when temperatures are below 32° F (usually nighttime lows). Apply to crops when temperatures have warmed above 35° F and crop is free of moisture of melting frost or heavy dew.
- DO NOT MIX with insecticides or fungicides incompatible with oil. Consult your State Agricultural Extension Service for specific compatibilities.
- DO NOT APPLY to trees weakened by disease, drought, drying winds, or high nitrogen applications.
- MAINTAIN CONSTANT VIGOROUS AGITATION during mixing and application to keep spray oil and water in solution.
- CONCENTRATE APPLICATIONS can delay bud development.

**CROPS: APPLES, ALMONDS, APRICOTS, CHERRIES, NECTARINES, OLIVES, PEACHES, PEAR, PLUMS, PRUNES, WALNUTS**

**PESTS:** San Jose Scale, Brown Apricot Scale, Italian Pear Scale, Parlatoria Scale, Oleander Scale, Pear Psylla, Aphids, Mites

**DORMANT OR DELAYED DORMANT APPLICATION**

Dilute: Apply 200 – 800 gallons of spray solution per acre using 1 – 1.5 gallons of oil per 100 gallons of water. For apples and pears, apply from 2-4 gallons of oil per 100 gallons of water.

**CONCENTRATE:** 5- 8 gallons per acre in minimum 50 gallons per acre

**FURTHER USE INSTRUCTIONS:** Do not apply to oil sensitive varieties due to fruit russetting or phytotoxicity to plant foliage or wood. Apply at the rate of 100 to 400 gallons of finished spray mix per acre. Walnuts – check with local farm advisor regarding all applications to walnuts.

**CROPS: ORNAMENTALS & SHADE TREES (Conifers, Holly, Pyracantha, Sycamore, Hawthorn, Flowering Peach, Flowering Cherry, Apple, Roses, Magnolia, Dogwood, Oak, Privet, Birch)**

**PESTS:** Black Scale, Brown Apricot Scale, Oyster Shell Scale, San Jose Scale, Soft Brown Scale, Red Spider Mites and Eggs, Aphids, Adelgids, Eriophyid Mites, Gall Mites, Lace Bug, Leaf Beetle larvae, Leafminer, Mealybug, Psyllids (immature), Sawfly, Whiteflies (immature)

**FOLIAR APPLICATION:** Apply 100 – 250 gallons of spray solution per acre using 0.5 – 1.0 gallons of oil per 100 gallons of water

**DORMANT OR DELAYED DORMANT APPLICATION:** 2 – 3 gallons

**FURTHER USE INSTRUCTIONS:** Conifer applications to Blue Spruce and Blue Juniper varieties may cause discoloration. May cause injury to Arborvitae and other cedars. Do not use on conifers or other ornamental plants not tolerant to oil sprays. Do not use dormant sprays on broadleaf evergreens such as Rhododendron, Cherry Laurel, Holly, Ivy, Azalea or Oregon Grape. In colder climates Cotoneaster, Barberry and Yews may be sensitive. Do not spray dormant sprays below 35° F. If handgun has not been used for twenty minutes or longer, recirculate back into tank until hose is flushed. Dormant Spray: Allow at least a 2-week interval between applications of dormant sprays. Do not spray dormant sprays when buds have fully formed, and shoot elongation is occurring. Do not spray dormant sprays when leaves are in obvious moisture deficit or plants are under stress. Reduce rate to 2 gallons/100 water for Fall dormant applications to only: American Red Oak, Japanese Black Pine, Dogwood, Weeping Cherry, Comelina Cherry, Crab Apple, Norway Maple, Pure Leaf Plum. Keep away from open blooms – spotting has been observed with oil use on open blooms of certain varieties.

**NOTICE**

INTEGRATED AGRIBUSINESS PROFESSIONALS (IAP) WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSE STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF IAP. TO THE EXTENT CONSISTENT WITH APPLICABLE LAWS, IN NO CASE SHALL IAP BE LIABLE FOR THE CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAWS, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAWS, EXCEPT AS EXPRESSLY PROVIDED HEREIN, IAP MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.

**Formulated for  
Integrated Agribusiness Professionals**

**7108 N. Fresno Street, Ste.150 • Fresno, CA 93720**

Integrated Agribusiness Professionals, IAP and the IAP logo are trademarks of Integrated Agribusiness Professionals Cooperative Inc.