



## IMPACT Herbicide

EPA REG. NO. 5481-524

### SUPPLEMENTAL LABEL

For control of emerged weeds  
in all corn types



### SUPPLEMENTAL LABEL

**Increased maximum single and annual use rate in corn to 2.0 fluid ounce (0.044 lb topramezone) per acre.**

EPA Reg. No.: 5481-524

#### Directions for Use

This supplemental label expires April 1, 2024 and must not be used or distributed after this date.

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.**

Before applying **IMPACT** please read the complete label affixed to the product container. All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Use of **IMPACT** according to this supplemental labeling is subject to the use precautions and limitations imposed by the label on the container.

**IMPORTANT:** Read the entire Directions for Use and the Limited Warranty and Disclaimer on the EPA-registered IMPACT herbicide label before using this product.

#### Use Information

For use in FIELD CORN (grown for grain, silage or seed), POPCORN (ear, kernel or seed) and SWEET CORN (ear, kernel or seed).

**Table 1. Emerged Broadleaf Weeds Controlled or Partially Controlled with IMPACT and Maximum Weed Size at Application**

Broadleaf Weeds <sup>1</sup>	IMPACT 0.75 fl oz per acre	IMPACT 1.0 fl oz per acre
	Weeds ≤ 4 inches in Size <sup>2</sup>	Weeds ≤ 6 inches in Size <sup>2</sup>
Amaranth, Palmer	C	C
Amaranth, Powell	C	C
Bindweed, field	PC	PC

Broadleaf Weeds <sup>1</sup>	IMPACT 0.75 fl oz per acre	IMPACT 1.0 fl oz per acre
	Weeds ≤ 4 inches in Size <sup>2</sup>	Weeds ≤ 6 inches in Size <sup>2</sup>
Burcucumber	C <sup>3,4</sup>	C <sup>4</sup>
Buckwheat, wild	NC	C <sup>3,4</sup>
Canola, volunteer	C	C
Carpetweed	C	C
Chickweed, common	C	C
Cocklebur, common	C	C
Dandelion	PC	PC
Devil's-claw	C <sup>3,4</sup>	C <sup>4</sup>
Galinsoga, hairy	C	C
Henbit	C	C
Jimsonweed	C	C
Kochia	C <sup>4</sup>	C <sup>4</sup>
Ladysthumb	C <sup>4</sup>	C <sup>4</sup>
Lambsquarters, common	C	C
Lettuce, prickly	C	C
Mallow, common	C <sup>3,4</sup>	C <sup>3,4</sup>
Mallow, Venice	C <sup>3,4</sup>	C <sup>3,4</sup>
Marestail (horseweed)	C	C
Morningglory, entireleaf	C <sup>3,4</sup>	C <sup>3,4</sup>
Morningglory, ivyleaf	C <sup>3,4</sup>	C <sup>3,4</sup>
Mustard, wild	C	C
Nightshade, black	C	C
Nightshade, Eastern black	C	C
Nightshade, hairy	C	C
Pigweed, prostrate	C	C
Pigweed, redroot	C	C
Pigweed, smooth	C	C
Pigweed, tumble	C	C
Purslane, common	PC <sup>3</sup>	C <sup>3,4</sup>
Pusley, Florida	C <sup>3</sup>	C
Ragweed, common	C	C
Ragweed, giant	C	C
Shepherd's-purse	C	C
Sida, prickly	C <sup>3,4</sup>	C <sup>4</sup>
Smartweed, Pennsylvania	C <sup>4</sup>	C <sup>4</sup>
Sunflower, wild (common)	C	C
Thistle, Canada	PC <sup>3</sup>	PC
Thistle, Russian	C	C
Velvetleaf	C	C
Waterhemp, common	C	C
Waterhemp, tall	C	C

<sup>1</sup>Refer to Section X for a list of scientific names on the EPA label.

<sup>2</sup>Broadleaf Weed Rating: C=Control; PC=Partial Control; NC=Not Controlled.

<sup>3</sup>Apply before weed exceeds 3 inches in height.

<sup>4</sup>The addition of atrazine at minimum 0.5 lb per acre is required for control.

**Table 2. Emerged Grass Weeds Controlled or Partially Controlled with IMPACT at 0.75 to 2.0 fl oz/A and Maximum Weed Size at Application**

Grass Weeds <sup>1</sup>	IMPACT 0.75 fl oz per acre	IMPACT 1.0 fl oz per acre	IMPACT 1.5 – 2.0 fl oz per acre
	Grasses ≤ 3 Inch Size <sup>2</sup>	Grasses ≤ 4 Inch Size <sup>2</sup>	Grasses ≤ 4 Inch Size <sup>2</sup>
Barnyardgrass	C <sup>5</sup>	C	C
Crabgrass, large	C <sup>5</sup>	C <sup>5</sup>	C
Crabgrass, smooth	C <sup>5</sup>	C <sup>5</sup>	C
Cupgrass, woolly	PC	C <sup>3</sup>	C
Foxtail, giant	C	C	C
Foxtail, green	C <sup>5</sup>	C	C
Foxtail, yellow	PC	C <sup>3,5</sup>	C
Goosegrass	C	C	C
Johnsongrass, rhizome	NC	NC	PC <sup>4</sup>
Johnsongrass, seedling	PC	C	C
Millet, wild-proso	C	C	C
Panicum, fall	PC	C <sup>3,5</sup>	C
Panicum, Texas	PC	PC	C <sup>4,5</sup>
Sandbur, field	NC	PC	C <sup>4,5</sup>
Shattercane	PC	PC	C <sup>4,5</sup>
Signalgrass, broadleaf	PC	C <sup>3,5</sup>	C <sup>5</sup>

<sup>1</sup>Refer to Section X for a list of scientific names on the EPA label.

<sup>2</sup>Grass Weed Rating: C=Control; PC=Partial Control; NC = Not Controlled.

<sup>3</sup>Apply before grass exceeds 3 inches in height.

<sup>4</sup>For best performance on these grass weeds use a rate of 2 fl oz per acre.

<sup>5</sup>The addition of atrazine at minimum 0.5 lb per acre is required for control.

### Restrictions

- Maximum use rate per year: Do not apply more than 0.044 lbs. of topramezone per acre or the equivalent of 2.0 fluid ounces of IMPACT per acre.
- DO NOT apply IMPACT within 45 days of corn harvest (fresh market sweet corn, silage, fodder, or grain).
- DO NOT graze or feed treated corn forage, silage, fodder, or grain for at least 45 days after an application of IMPACT.
- DO NOT apply this product through any type of irrigation system.

©2020 AMVAC Chemical Corporation is a wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. AMVAC, Impact, and their respective logos are trademarks owned by AMVAC Chemical Corporation.

Registrant:

AMVAC Chemical Corporation  
4695 MacArthur Court, Suite 1200  
Newport Beach, CA 92660 U.S.A.