



**FIFRA 24(c) SPECIAL LOCAL NEED LABEL (SLN)
FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA**

For Aerial Application to Almonds, Stone Fruit Crop Group 12-12 with Standing Water

Inspire Super®

EPA Reg. No. 100-1317

EPA SLN No. CA-210002

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2025.

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA-registered Inspire Super, EPA Reg No. 100-1317 label and this label.

Crop	Diseases	Product Rate fl oz/Acre	Remarks
Almonds	Alternaria Leaf Spot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum acutatum</i>) Blossom Blight (<i>Monilinia</i> spp.) Green Fruit Rot (<i>Botrytis cinerea</i>) Leaf Blight (<i>Seimatosporium lichenicola</i>) Leaf Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophila</i>) Shot Hole (<i>Wilsonmyces carpophilus</i>)	16 - 20	For blossom blight, apply 16-20 fl oz of Inspire Super during the bloom period. For Alternaria leaf spot and scab, begin applications prior to disease onset when conditions are conducive for disease. If monitoring or history indicates the presence of Alternaria, apply 20 fl oz/A of Inspire Super in the late spring (mid-April to beginning of May) and then repeat the treatment 2-3 weeks later. For all other diseases, use 16-20 fl oz/A. Begin applications prior to disease onset when conditions are conducive for disease. Apply Inspire Super on a 14- to 21-day schedule. Make no more than 2 sequential applications before alternating to another fungicide with a different mode of action. The minimum retreatment interval is 14 days. If disease pressure is high, use the shortest interval and highest rate.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Use a minimum of 10 gal/A of water for aerial applications. Use ground application for best results.

Specific Use Restrictions:

- 1) Make no more than two applications by air per year.
- 2) Do not apply more than 40 fl oz/A (0.65 lb ai cyprodinil/A) of Inspire Super per year.
- 3) Do not apply more than 0.46 lb ai/A/year per crop of difenoconazole-containing products.
- 4) Do not apply more than 1.4 lb ai/A/year of cyprodinil-containing products for almonds.
- 5) Do not apply within 60 days of harvest (60-day PHI).

Crop	Diseases	Product Rate fl oz/Acre	Remarks
Stone Fruit Crop Group 12-12* Apricots Cherries, Tart Nectarines Peaches Plums Plumcot Prunes And cultivars and/or hybrids of these.	Alternaria Spot and Fruit Rot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum</i> spp.) Brown Rot Blossom Blight and Fruit Rot <i>Monilinia fructicola</i> , <i>M. laxa</i>) Leaf Rust (<i>Tranzschelia discolor</i>) Powdery Mildew (<i>Sphaerotheca pannosa</i> , <i>Podosphaera clandestina</i>) Scab (<i>Cladosporium carpophilum</i>) Shot Hole (<i>Wilsonomyces carpophilus</i>)	16 - 20	For brown rot blossom blight, begin applications at early bloom and make a second application at full bloom. For brown rot on fruit, apply as needed a maximum of two sprays during the preharvest period up to the day of harvest (minimum of a 7-day retreatment interval). If high inoculum and severe disease conditions persist, apply a registered non-Group 3 fungicide. If disease pressure is high, use the highest rate.

***Stone Fruit Crop Group:** Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Use a minimum of 10 gal/A spray volume by air.

Specific Use Restrictions:

- 1) Make no more than two applications by air.
- 2) Do not apply more than 80 fl oz/A (1.3 lb ai cyprodinil/A) of Inspire Super per year.
- 3) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 4) Do not apply more than 1.4 lb ai/A/year of cyprodinil-containing products for Stone Fruit crop group 12-12.
- 5) Do not apply within 2 days of harvest (2-day PHI).
- 6) Do not apply more than a maximum total of 4 applications (air plus ground) per year.

Requirements:

1. For applications to orchards that have standing water due to a storm event.
2. Water must be standing and will not flow into state waterways where fish and wildlife are likely to be affected.
3. Applications are made in accordance with all other label directions.
4. The County Agriculture Commissioner must assure the required conditions are met.

Restrictions:

1. No fungicide application shall occur if:
 - a. Soil moisture is at field capacity and a storm event, forecasted by the National Oceanic and Atmospheric Administration (NOAA) or National Weather Service (NWS), is to occur within 48 hours following the application; or
 - b. A storm event likely to produce runoff from the treated area is forecasted by NOAA/NWS to occur within 48 hours following the application.
 2. Applications are not allowed if water is flowing off the intended application site.
 3. Standing water will not be pumped or removed from the orchard after the fungicide application.
 4. Do not apply this product through any type of irrigation system.
-

Inspire Super® registered trademark of a Syngenta Group Company

©2019 Syngenta

24(c) Registrant:
Syngenta Crop Protection, LLC.
P. O. Box 18300
Greensboro, North Carolina 27419
Telephone: 1-800-334-9481

Label Code: CA1317190AA0819