

# Supplemental Labeling



Dow AgroSciences LLC      9330 Zionsville Road      Indianapolis, IN 46268-1054 USA

## Intrepid<sup>®</sup> 2F EPA Reg. No. 62719-442

### Control of Lepidoptera Larvae Infesting Root Vegetables (Subgroups 1A, 1B)<sup>1</sup> (Not for Use and Distribution in New York State)

**ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid<sup>®</sup> 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

#### Directions for Use

<sup>1</sup>Root vegetables (subgroups 1A, 1B) including black salsify, carrot, celeriac, chicory, edible burdock, garden beet, ginseng, horseradish, parsnip, oriental radish, radish, rutabaga, salsify, skirret, Spanish salsify, sugarbeet, turnip, turnip-rooted chervil, and turnip-rooted parsley

Refer to product label for Product Information, Mixing Directions and Application instructions.

| Pests  | Application Rate<br>(fl oz/acre)      | Application Timing  | Restrictions  |
|--|---------------------------------------|---|---|
| armyworms<br>cabbageworms<br>cutworm<br>(suppression only)<br>loopers<br>saltmarsh caterpillar<br>webworms | 8 - 16<br>(0.12 – 0.25 lb<br>ai/acre) | Apply at egg hatch or when first signs of feeding occur. Use a higher rate for heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapply to protect new growth until moth flights and/or hits subside. | <ul style="list-style-type: none"> <li>• <b>Preharvest Interval:</b> Do not apply within 1 day of harvest for all root vegetables except sugar beet. Do not apply within 7 days of sugarbeet harvest.</li> <li>• Do not apply more than a total of 64 fl oz of Intrepid 2F (1 lb ai) per acre per year for all crops except radish.</li> <li>• Do not apply more than a total of 32 fl oz of Intrepid 2F (0.5 lb ai) per acre per year for radish.</li> </ul> |

**Ground Application:** Apply in a minimum of 10 gallons per acre by conventional ground equipment. Use a spray volume that assures uniform coverage of the infested portions of the treated crop. Calibrate equipment to the desired spray volume.

**Aerial Application:** Apply in a minimum of 10 gpa. Use a higher carrier volume for heavy infestations and in situations where thorough coverage is difficult to achieve.

**Spray Adjuvants:** The addition of agricultural adjuvants to sprays of Intrepid 2F may improve initial spray deposits, redistribution and weatherability. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Always add adjuvants last in the mixing process.

**Note:** Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R846-073

EPA accepted 7/11/12

Initial printing.