

# Supplemental Labeling



Dow AgroSciences LLC      9330 Zionsville Road      Indianapolis, IN 46268-1054 USA

## Intrepid<sup>®</sup> 2F EPA Reg. No. 62719-442

### Control of Lepidoptera Larvae Infesting Tropical Tree Fruits<sup>1</sup> (Not for Use and Distribution in New York State)

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid<sup>®</sup> 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

#### Directions for Use

<sup>1</sup>Tropical tree fruits including acerola, atemoya, avocado, biriba, black sapote, canistel, cherimoya, custard apple, feijoa, guava, ilama, jaboticaba, longan, lychee, mamey sapote, mango, papaya, passionfruit, pulasan, rambutan, sapodilla, soursop, Spanish lime, star apple, starfruit, sugar apple, wax jambu

Refer to product label for General Information, Mixing and Application directions.

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
lepidopteran larvae including European grapevine moth guava moth ( <i>Argyresthia</i> ) leafrollers light brown apple moth loopers orange tortrix spanworms webbing worms western tussock moth	10 - 16 (0.16 - 0.25 lb ai/acre)	Apply at egg hatch or when first signs of feeding occur. Use a higher rate for heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapply at a 6- to 10-day re-treatment interval to protect new growth until moth flights and/or hits subside.	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 64 fl oz of Intrepid 2F (1 lb ai) per acre per year.</li> <li>• Do not make more than 5 applications per calendar year.</li> <li>• <b>Acerola, Feijoa, Guava, Jaboticaba, Passionfruit, Starfruit, Wax Jambu</b>  <b>Preharvest Interval:</b> Do not apply within 3 days of harvest.  <b>Minimum Re-treatment Interval:</b> 6 days</li> </ul>

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
			<ul style="list-style-type: none"> <li data-bbox="1101 178 1425 451"> <b>• Atemoya, Avocado, Biriba, Cherimoya, Custard Apple, Ilima, Soursop, Sugar Apple</b>  <b>Preharvest Interval:</b> Do not apply within 2 days of harvest.  <b>Minimum Re-treatment Interval:</b> 6 days </li> <li data-bbox="1101 489 1425 762"> <b>• Black Sapote, Canistel, Mamey Sapote, Mango, Papaya, Sapodilla, Star Apple</b>  <b>Preharvest Interval:</b> Do not apply within 3 days of harvest.  <b>Minimum Re-treatment Interval:</b> 10 days </li> <li data-bbox="1101 800 1425 978"> <b>• Longan, Lychee, Pulasan, Rambutan, Spanish Lime</b>  <b>Preharvest Interval:</b> Do not apply within 14 days of harvest. </li> <li data-bbox="1101 989 1425 1041"> <b>• Minimum Re-treatment Interval:</b> 10 days </li> </ul>

**Ground Application:** Apply in a minimum of 50 gpa by conventional ground equipment to trees 10 feet tall or less. For trees greater than 10 feet tall, apply in a minimum of 100 gpa by conventional group equipment. Use a higher carrier volume for heavy infestations and in situations where thorough coverage is difficult to achieve. Use a spray volume that assures uniform coverage of the infested portions of the treated crop. Calibrate equipment to the desired spray volume.

**Aerial Application:** Apply in a minimum of 10 gpa. Use a higher carrier volume for heavy infestations and in situations where thorough coverage is difficult to achieve.

**Spray Adjuvants:** The addition of agricultural adjuvants to sprays of Intrepid 2F may improve initial spray deposits, redistribution and weatherability. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Always add adjuvants last in the mixing process.

**Note:** Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R846-090

EPA accepted 10/29/13

Initial printing.