

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid Edge[®]

EPA Reg. No. 62719-666

Special 2(ee) Recommendation[†] For Control of Thrips in Cotton

(For distribution and use in AL, AR, LA, MS, MO, and TN only)

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Intrepid Edge[®] insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Intrepid Edge according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid Edge.

Directions for Use: Refer to the Intrepid Edge[®] product label for Application Directions, Mixing Directions, and Restrictions.

Application Rate: The amount of Intrepid Edge[®] applied per acre will depend upon pest infestation level. Use a lower rate in the rate range for light infestations and a higher rate in the rate range for heavy infestations. Use of an adjuvant may increase the level of thrips control.

Application Timing: Make applications during late cotyledon stage cotton and then continue to manage populations until cotton is no longer susceptible to early season thrips damage. Under heavy pressure multiple applications may be needed to control pest. Using Intrepid Edge[®] is not a replacement for insecticide seed treatments or in-furrow insecticide applications at planting. Early season, foliar applications of Intrepid Edge[®] to suppress thrips in cotton should only be made to fields that have already received an insecticide seed treatment or in-furrow insecticide application for thrips control.

Resistance Management: Do not make more than two consecutive applications of Group 5 or 18 insecticides per crop. If additional treatments are required after two consecutive applications of Group 5 or 18 insecticides, rotate to another class of effective insecticide for at least one application. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Avoid applying Intrepid Edge and other Group 5 and 18 insecticides to consecutive generations of all target insect pests.

Pest	Intrepid Edge [®] (fl oz/acre)
Thrips	3 – 6

Restrictions:

- **Preharvest Interval:** Do not apply within 28 days of harvest.
- Do not apply more than 8 fl oz of Intrepid Edge[®] per acre per application.
- Do not apply more than a total of 48 fl oz of Intrepid Edge[®] (0.94 lb ai methoxyfenozide, 0.19 lb ai spinetoram) per acre per year.
- **Minimum Treatment Interval:** Do not make applications less than 4 days apart.
- **Maximum Number of Applications:** Do not make more than 6 applications per year.
- Do not make more than two consecutive applications.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow
R911-011
Issued: 03/20/18
Initial printing.