

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid Edge[®]

EPA Reg. No. 62719-666

Special 2(ee) Recommendation[†]

For Control of Citrus Leafminer and Asian Citrus Psyllid in Citrus in Florida

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Intrepid Edge[®] insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Intrepid Edge according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid Edge.

Directions for Use

Pest	Intrepid Edge [®] (fl oz/acre)
Citrus leafminer Asian Citrus Psyllid ¹	8 – 10

¹Residual control of Asian Citrus Psyllid may not be expected at these rates. Continue to monitor infestations with regularity.

Refer to the Intrepid Edge[®] product label for General Use Precautions, Mixing Directions, Application Directions, and Restrictions.

Application Timing: Treat in accordance with local economic thresholds. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Resistance Management: Do not make more than two consecutive applications of Group 5 or 18 insecticides per crop. If additional treatments are required after two consecutive applications of Group 5 or 18 insecticides, rotate to another class of effective insecticide for at least one application. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Avoid applying Intrepid Edge and other Group 5 and 18 insecticides to consecutive generations of all target insect pests.

Restrictions:

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- Do not apply more than 12 fl oz of Intrepid Edge[®] per acre per application.
- Do not apply more than a total of 36 fl oz of Intrepid Edge[®] (0.70 lb ai methoxyfenozide, 0.14 lb ai spinetoram) per acre per season.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- **Maximum Number of Applications:** Do not make more than 3 applications per year.
- Do not apply in citrus nurseries or citrus in greenhouses.

[®]Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R911-009

Issued: 01/30/18

Replaces R911-008