

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid Edge[®]

EPA Reg. No. 62719-666

Special 2(ee) Recommendation[†]

For Control of Walnut Husk Fly in Walnut

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Intrepid Edge[®] insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Intrepid Edge according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid Edge.

Directions for Use: Refer to the Intrepid Edge[®] product label for General Use Precautions, Mixing Directions, Application Directions, and Restrictions.

Application Rate: The amount of Intrepid Edge[®] applied per acre will depend upon tree size and pest infestation level. Use a lower rate in the rate range for light infestations and/or smaller trees and a higher rate in the rate range for heavy infestations and/or larger trees.

Ground Application: Apply in a minimum of 50 gpa by conventional ground equipment. For best results, use 100 to 200 gpa. Use a spray volume that assures uniform coverage of the infested crop.

Aerial Application: Apply in a minimum of 10 gpa. This method may result in reduced efficacy if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

Resistance Management: Do not make more than two consecutive applications of Group 5 or 18 insecticides per crop. If additional treatments are required after two consecutive applications of Group 5 or 18 insecticides, rotate to another class of effective insecticide for at least one application. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Avoid applying Intrepid Edge and other Group 5 and 18 insecticides to consecutive generations of all target insect pests.

Pest	Intrepid Edge [®] (fl oz/acre)
Walnut husk fly	12 – 14

Restrictions:

- **Preharvest Interval:** Do not apply within 14 days of harvest.
- Do not apply more than 18 fl oz of Intrepid Edge[®] per acre per application.
- Do not apply more than a total of 51 fl oz of Intrepid Edge[®] (1 lb ai methoxyfenozide, 0.2 lb ai spinetoram) per acre per season.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- **Maximum Number of Applications:** Do not make more than 4 applications per year.
- Do not make more than two consecutive applications.