

Tree Nuts Crop Grouping: Almond, Areca (Betelnut Palm), Beechnut, Chestnut, Chinquapin, Filbert, Hickory nut, Kiwi, Macadamia nut, Pecan, Pistachio and Walnut.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. **Do not double spray or reduce ground speed of spraying device which would cause label rates to be exceeded. Avoid applying tank mix overage onto the crop which would exceed labeled rates.** Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. **During periods of excessive temperatures use lower product label rates.** Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).